arXiv:0810.5165v2 [math.RT] 22 Jul 2009

# SOME COMPUTATIONS ABOUT KAZHDAN-LUSZTIG CELLS IN AFFINE WEYL GROUPS OF RANK 2

JÉRÉMIE GUILHOT

Abstract. In the last section of the paper "Generalized induction of Kazhdan-Lusztig cells" and in "Kazhdan-Lusztig cells in affine Weyl groups of rank 2" the author described the partition into Kazhdan-Lusztig cells of the affine Weyl groups of rank 2 for all choices of parameters. The proof of these results relies on some explicit computation with GAP [4]. In these notes we give some detail of these computations.

## 1. Introduction

These notes are organized as follows:

- in Section 2, we give some details in the computations involved in the last section of [2, §6];
- in Section 3, we give some details in the computations involved in the proof of [3, Theorem 5.1] for the affine Weyl group of type $\tilde{G}_2$. We also give the left order on the left cells for all choices of parameters.
- in Section 4, we describe the partition of the affine Weyl group of type $\tilde{B}_2$ into cells for all parameters.
- in Section 5, we give some details in the computations involved in the proof of [3, Theorem 5.3] for the affine Weyl group of type $\tilde{B}_2$. We also give the left order on the left cells for all choices of parameters.

## 2. Computations in [2, §6]

Let $W$ be the affine Weyl group of type $\tilde{G}_2$ with diagram and weight function given by



where $a, b$ are positive integers such that $a/b > 4$. In this section we go through the proof of Lemma 6.4, Claim 6.6 and Claim 6.8 in [2] and give some details in the computations involved.

We keep the notation of [2, §6]. Recall that



Figure 1. Decomposition of $\tilde{G}_2$ into left cells in the case $a > 4b$

### 2.1. **The sets $C_i$.**

For $1 \leq i \leq 6$, let

1. $u_i \in C_i$ be the element of minimal length in $C_i$;
2. $v_i \in A_i$ be the element of minimal length in $A_i$;
3. $v'_i \in A'_i$ be the element of minimal length in $A'_i$.

We set $U := \{u_i, v_i, v'_i \mid 1 \leq i \leq 6\}$, $X_{v_i} = X_{v'_i} = X_{\{s_1,s_2\}}$ and

$$X_{u_i} = \{z \in W \mid zu_i \in C_i\}.$$

In this section we prove the following.

1. The submodule
$$\mathcal{M} := \langle T_x C_u \mid u \in U, x \in X_u \rangle_{\mathcal{A}} \subseteq \mathcal{H}$$
is a left ideal.

(2) Let $u \in U$, $u' < u$ and $x \in X_u$. We have

$$P_{u',u}T_xC_u \equiv T_{xu} \mod \mathcal{H}_{<0}$$

First we prove (1). To this end we only need to show that $T_{s_2}C_{u_i}$ and $T_{s_2s_1s_2s_1s_2s_3}C_{u_i}$ lie in $\mathcal{M}$ for all $1 \leq i \leq 6$. We have

$$T_{s_2}C_{u_i} = C_{v_i} - v^{-b}C_{u_i} \quad \text{for } i = 1,2,3,6$$

and

$$\begin{aligned} T_{s_2}C_{u_4} &= C_{v_4} - v^{-b}C_{u_4} + C_{v_2} \\ T_{s_2}C_{u_5} &= C_{v_5} - v^{-b}C_{u_5} + C_{v_1}. \end{aligned}$$

Thus $T_{s_2}C_{u_i} \in \mathcal{M}$.
For $i = 1,2,3,6$ we have

$$\begin{aligned} T_3C_{u_i} &= C_{3u_i} + \mathcal{A}C_{u_i} \\ T_{23}C_{u_i} &= C_{23u_i} + \mathcal{A}C_{3u_i} + \mathcal{A}C_{u_i} + \mathcal{A}C_{v_i} \\ T_{123}C_{u_i} &= C_{123u_i} + \mathcal{A}C_{23u_i} + \mathcal{A}C_{3u_i} + \mathcal{A}C_{u_i} + \mathcal{A}C_{v_i} \\ T_{2123}C_{u_i} &= C_{2123u_i} + \mathcal{A}C_{123u_i} + \mathcal{A}C_{3u_i} + \mathcal{A}C_{u_i} \\ T_{12123}C_{u_i} &= C_{12123u_i} + \mathcal{A}C_{2123u_i} + \mathcal{A}C_{123u_i} + \mathcal{A}C_{u_i} \\ T_{212123}C_{u_i} &= C_{v_i'} + \mathcal{A}C_{12123u_i} + \mathcal{A}C_{123u_i} + \mathcal{A}C_{u_i} + \mathcal{A}C_{v_i} \end{aligned}$$

thus

$$C_{3u_i}, C_{23u_i}, C_{123u_i}, C_{2123u_i}, C_{12123u_i} \in \mathcal{M}.$$

Since $C_{v_i'}, C_{u_i}, C_{v_i} \in \mathcal{M}$ we see that $T_{212123}C_{u_i} \in \mathcal{M}$ as required.
We have

$$\begin{aligned} T_3C_{u_4} &= C_{3u_4} + \mathcal{A}C_{u_4} \\ T_{23}C_{u_4} &= C_{23u_4} + \mathcal{A}C_{3u_4} + \mathcal{A}C_{u_4} + \mathcal{A}C_{v_4} + \mathcal{A}C_{v_2} \\ T_{123}C_{u_4} &= C_{123u_4} + \mathcal{A}C_{23u_4} + \mathcal{A}C_{3u_4} + \mathcal{A}C_{u_4} + \mathcal{A}C_{v_4} + \mathcal{A}C_{v_2} \\ T_{2123}C_{u_4} &= C_{2123u_4} + \mathcal{A}C_{123u_4} + \mathcal{A}C_{3u_4} + \mathcal{A}C_{u_4} \\ T_{12123}C_{u_4} &= C_{12123u_4} + \mathcal{A}C_{2123u_4} + \mathcal{A}C_{123u_4} + \mathcal{A}C_{u_4} \\ T_{212123}C_{u_4} &= C_{v_i'} + \mathcal{A}C_{12123u_4} + \mathcal{A}C_{123u_4} + \mathcal{A}C_{u_4} + \mathcal{A}C_{v_4} + \mathcal{A}C_{v_2} \end{aligned}$$

thus

$$C_{3u_i}, C_{23u_i}, C_{123u_i}, C_{2123u_i}, C_{12123u_i} \in \mathcal{M}.$$

Since $C_{v_i'}, C_{u_i}, C_{v_i} \in \mathcal{M}$ we see that $T_{212123}C_{u_4} \in \mathcal{M}$ as required.
We have

$$\begin{aligned} T_3C_{u_5} &= C_{3u_5} + \mathcal{A}C_{u_5} \\ T_{23}C_{u_5} &= C_{23u_5} + \mathcal{A}C_{3u_5} + \mathcal{A}C_{u_5} + \mathcal{A}C_{v_5} + \mathcal{A}C_{v_1} \\ T_{123}C_{u_5} &= C_{123u_5} + \mathcal{A}C_{23u_5} + \mathcal{A}C_{3u_5} + \mathcal{A}C_{u_5} + \mathcal{A}C_{v_5} + \mathcal{A}C_{v_1} \\ T_{2123}C_{u_5} &= C_{2123u_5} + \mathcal{A}C_{123u_5} + \mathcal{A}C_{3u_5} + \mathcal{A}C_{u_5} \\ T_{12123}C_{u_5} &= C_{12123u_5} + \mathcal{A}C_{2123u_5} + \mathcal{A}C_{123u_5} + \mathcal{A}C_{u_5} \\ T_{212123}C_{u_5} &= C_{v_i'} + \mathcal{A}C_{12123u_5} + \mathcal{A}C_{123u_5} + \mathcal{A}C_{u_5} + \mathcal{A}C_{v_5} + \mathcal{A}C_{v_1} \end{aligned}$$

thus

$$C_{3u_i}, C_{23u_i}, C_{123u_i}, C_{2123u_i}, C_{12123u_i} \in \mathcal{M}.$$

Since $C_{v'_i}, C_{u_i}, C_{v_i} \in \mathcal{M}$ we see that $T_{212123}C_{u_5} \in \mathcal{M}$ as required.
Statement (1) follows.

Before proving (2), we need to introduce some definitions. For $w \in W$ we set

$$Y(w) = \{y \in W | \ell(wy^{-1}) = \ell(w) - \ell(y)\}.$$

We have $y \in Y(w)$ if and only if there exists a reduced expression $s_1 \dots s_n$ of $w$ and $1 \leq i \leq n$ such that $y = s_i \dots s_n$. Recall the definition of translation in [1, Definition 4.1].

**Claim 2.1.** Let $w \in W$ be a translation. Let $z \in W$. The following statements are equivalent

(1) $\ell(wz) = \ell(w) + \ell(z)$;
(2) $\ell(xz) = \ell(x) + \ell(z)$ for all $x \in \{zw^n | z \in Y(w), n \in \mathbb{N}\}$.

*Proof.* The fact that (2) implies (1) is clear since $w \in \{zw^n | z \in Y(w), n \in \mathbb{N}\}$. Assume that (1) holds. Then using [1, Lemma 4.2] in its right version we get

$$\ell(wz) = \ell(w) + \ell(z) \Leftrightarrow \ell(w^n z) = \ell(w^n) + \ell(z).$$

Let $x \in \{zw^n | z \in Y(w), n \in \mathbb{N}\}$. Let $y \in Y_w$ and $n \in \mathbb{N}$ be such that $x = yw^n$. We have

$$\ell(wz) = \ell(w) + \ell(z) \Rightarrow \ell(w^{n+1}z) = \ell(w^{n+1}) + \ell(z) \Rightarrow \ell(yw^n z) = \ell(yw^n) + \ell(z)$$

as required. □

We now prove (2). Let $w = s_1s_2s_1s_2s_3s_1s_2s_1s_2s_3$. Note that $w$ is a translation. We have

$$X_{u_i} = \{yw^n | y \in Y(w), n \in \mathbb{N}\}.$$

Let $u < u_i$. Using GAP one can show that for all $y \in Y(w)$ we have

$$P_{u,u_i}T_yC_u \equiv T_{yu} \mod \mathcal{H}_{<0}.$$

Next, one can show that

$$P_{u,u_i}T_wC_u = T_{wu} + \sum_{z \in W} a_z T_z$$

where $a_z \in \mathcal{A}_{<0}$ and if $a_z \neq 0$ then $z$ satisfy $\ell(wz) = \ell(w) + \ell(z)$. Now let $x = yw^n \in X_{u_i}$ (where $y \in Y(w)$ and $n \in \mathbb{N}^*$). Then using Claim 2.1 we obtain

$$\begin{aligned} P_{u,u_i}T_xC_u &= T_{yw^{n-1}}\Big(T_{wu} + \textstyle\sum_{z \in W} a_z T_z\Big) \\ &= T_{xu} + \sum_{z \in W} a_z T_{yw^{n-1}z} \\ &\equiv T_{xu} \mod \mathcal{H}_{<0} \end{aligned}$$

and statement (2) follows.

## 2.2. **Proof of Claim 6.6.**

Let $u = s_1s_3s_2s_1$ and recall that

$$\begin{aligned} u_1 &= s_1s_2s_1s_2s_1,\\ v_1 &= s_1s_2s_1s_2s_1s_2,\\ v_1' &= s_1s_2s_1s_2s_1s_2s_3s_2s_1s_2s_1,\\ u_2 &= s_1s_2s_1s_2s_1s_3s_2s_1,\\ v_2 &= s_2s_1s_2s_1s_2s_1s_3s_2s_1,\\ v_2' &= s_2s_1s_2s_1s_2s_3s_1s_2s_1s_2s_1s_3s_2s_1\\ v_3 &= s_2s_1s_2s_1s_2s_1s_3, \end{aligned}$$

and

$$\begin{aligned} X_u &:= \{z \in W | zu \in B_1\}\\ X_{u_i} &= \{z \in W \mid zu_i \in C_i\},\\ X_{v_i} &= X_{v_i'} = X_{\{s_1,s_2\}} \quad \text{for } 1 \leq i \leq 6. \end{aligned}$$

Let $U := \{u, u_1, v_1, v_1', u_2, v_2, v_2', v_3\}$. In this section we prove the following.

1. The submodule
$$\mathcal{M} := \langle T_xC_v \mid v \in U, x \in X_u\rangle_{\mathcal{A}} \subseteq \mathcal{H}$$
is a left ideal.
2. Let $u' < u$ and $x \in X_u$. We have
$$P_{u',u}T_xC_u \equiv T_{xu} \mod \mathcal{H}_{<0}$$

We prove (1). To this end it is enough to show that $T_{1212}C_u \in \mathcal{M}$ and $T_{1212132}C_u$ lie in $\mathcal{M}$. We have

$$\begin{array}{rcl} T_2C_u &=& C_{2u} + \mathcal{A}C_u\\ T_{12}C_u &=& C_{12u} + \mathcal{A}C_{2u} + \mathcal{A}C_u\\ T_{212}C_u &=& C_{212u} + \mathcal{A}C_{12u} + \mathcal{A}C_u\\ T_{1212}C_u &=& C_{u_2} + \mathcal{A}C_{212u} + \mathcal{A}C_{12u} + \mathcal{A}C_{u_1} + \mathcal{A}C_{v_1} \end{array}$$

Thus $T_{1212}C_u \in \mathcal{M}$.

We have

$$\begin{array}{rcl}
T_2C_u & = & C_{2u}+\mathcal{A}C_u \\
T_{32}C_u & = & C_{32u}+\mathcal{A}C_{2u} \\
T_{132}C_u & = & C_{132u}+\mathcal{A}C_{32u}+\mathcal{A}C_{12u}+\mathcal{A}C_{2u}+\mathcal{A}C_u \\
T_{2132}C_u & = & C_{2132u}+\mathcal{A}C_{212u}+\mathcal{A}C_{132u}+\mathcal{A}C_{32u}+\mathcal{A}C_{12u}+\mathcal{A}C_{2u}+\mathcal{A}C_u \\
T_{12132}C_u & = & C_{12132u}+\mathcal{A}C_{2132u}+\mathcal{A}C_{212u}+\mathcal{A}C_{32u}+\mathcal{A}C_{12u}+\mathcal{A}C_{2u} \\
 & & +\mathcal{A}C_u+\mathcal{A}C_{u_1}+\mathcal{A}C_{v_1}+\mathcal{A}C_{u_2} \\
T_{212132}C_u & = & C_{212132u}+\mathcal{A}C_{12132u}+\mathcal{A}C_{2132u}+\mathcal{A}C_{212u}+\mathcal{A}C_{32u} \\
 & & +\mathcal{A}C_{12u}+\mathcal{A}C_{2u}+\mathcal{A}C_u+\mathcal{A}C_{u_1}+\mathcal{A}C_{v_2}+\mathcal{A}C_{u_2} \\
T_{1212132}C_u & = & C_{v'_1}+\mathcal{A}C_{212132u}+\mathcal{A}C_{2132u}+\mathcal{A}C_{212u}+\mathcal{A}C_{32u}+\mathcal{A}C_{2u} \\
 & & +\mathcal{A}C_{v_1}+\mathcal{A}C_{v_2}+\mathcal{A}C_{v_3}
\end{array}$$

Thus $T_{1212132}C_u \in \mathcal{M}$ as desired.

We prove (2). Let $w_1 = s_1s_2s_3s_2s_1s_2$ and $w_2 = s_2s_3s_2s_1s_2s_1$. These are both translations. Any $x \in X_u$ can be written under the form $yw_1^n$ with $y \in Y(w_1)$ or $yw_2^ns_3s_2$ with $y \in Y(w_2)$. One can show that for all $y_1 \in Y(w_1)$ and all $y_2 \in Y(w_2)$ we have

$$P_{u,u_i}T_{y_1}C_u \equiv T_{y_1u} \mod \mathcal{H}_{<0} \text{ and } P_{u,u_i}T_{y_2s_3s_2}C_u \equiv T_{y_2s_3s_2u} \mod \mathcal{H}_{<0}$$

Furthermore, we have

$$P_{u,u_i}T_{w_1}C_u = T_{w_1u} + \sum_{z\in W} a_zT_z$$

where $a_z \in \mathcal{A}_{<0}$ and if $a_z \neq 0$ then $z$ satisfy $\ell(w_1z) = \ell(w_1) + \ell(z)$. We have

$$P_{u,u_i}T_{w_2s_3s_2}C_u = T_{w_2s_3s_2u} + \sum_{z\in W} a'_zT_z$$

where $a'_z \in \mathcal{A}_{<0}$ and if $a'_z \neq 0$ then $z$ satisfy $\ell(w_2z) = \ell(w_2) + \ell(z)$. Arguing as in the previous section we obtain that (2) holds.

### 2.3. **Proof of Claim 6.8.**

Recall that $v = s_1s_3s_2s_1s_2s_3$ and

$$X_v := \{z \in W | zv \in B_3\} \qquad Y_v := \{x \in X_v | \ell(xs_2s_1s_2) = \ell(x) - 3\}.$$

Let $U = \{v, u_4, v_4, v'_4, v_3, v_2, v_5\}$. In this section we prove the following.

(1) The submodule

$$\mathcal{M} := \langle T_xC_u \mid u \in U, x \in X_u\rangle_{\mathcal{A}} \subseteq \mathcal{H}$$

is a left ideal.

(2) Let $x \in X_v - Y_v$. Then we have

$$T_xC_v \equiv T_{xv} \mod \mathcal{H}_{<0}$$

(3) Let $y \in Y_v$ and $y_0 = s_2s_1s_2$. We have

$$\tilde{C}_{yv} \equiv T_{yv} + T_{y_0v_3} \mod \mathcal{H}_{<0}$$

We prove (1). To this end it is enough to show that $T_{1212}C_u \in \mathcal{M}$ and $T_{1212132}C_u$ lie in $\mathcal{M}$. We have

$$\begin{array}{rcl}
T_2C_v & = & C_{2v} + \mathcal{A}C_v \\
T_{12}C_v & = & C_{12v} + \mathcal{A}C_{2v} + \mathcal{A}C_v \\
T_{212}C_v & = & C_{212v} + \mathcal{A}C_{12v} + \mathcal{A}C_v + \mathcal{A}C_{v_3} \\
T_{1212}C_v & = & C_{u_4} + \mathcal{A}C_{212v} + \mathcal{A}C_{12v} + \mathcal{A}C_{v_3}
\end{array}$$

Thus $T_{1212}C_v \in \mathcal{M}$. We have

$$\begin{array}{rcl}
T_2C_v & = & C_{2v} + \mathcal{A}C_v \\
T_{32}C_v & = & C_{32v} + \mathcal{A}C_{2v} \\
T_{132}C_v & = & C_{132v} + \mathcal{A}C_{32v} + \mathcal{A}C_{12v} + \mathcal{A}C_{2v} + \mathcal{A}C_v \\
T_{2132}C_v & = & C_{2132v} + \mathcal{A}C_{212v} + \mathcal{A}C_{132v} + \mathcal{A}C_{32v} + \mathcal{A}C_{12v} + \mathcal{A}C_{2v} + \mathcal{A}C_v + \mathcal{A}C_{v_3} \\
T_{12132}C_v & = & C_{12132v} + \mathcal{A}C_{2132v} + \mathcal{A}C_{212v} + \mathcal{A}C_{32v} + \mathcal{A}C_{12v} + \mathcal{A}C_{2v} \\
 & & +\mathcal{A}C_v + \mathcal{A}C_{u_4} + \mathcal{A}C_{v_3} \\
T_{212132}C_v & = & C_{212132v} + \mathcal{A}C_{12132v} + \mathcal{A}C_{2132v} + \mathcal{A}C_{212v} + \mathcal{A}C_{32v} \\
 & & +\mathcal{A}C_{12v} + \mathcal{A}C_{2v} + \mathcal{A}C_v + \mathcal{A}C_{u_4} + \mathcal{A}C_{v_4} + \mathcal{A}C_{v_3} + \mathcal{A}C_{v_2} \\
T_{1212132}C_v & = & T_{212123}C_{v_3} + \mathcal{A}C_{212132v} + \mathcal{A}C_{2132} + \mathcal{A}C_{212v} + \mathcal{A}C_{32v} + \mathcal{A}C_{2v} \\
 & & +\mathcal{A}C_{v_5} + \mathcal{A}C_{v_4} + \mathcal{A}C_{v_3} + \mathcal{A}C_{v_2}
\end{array}$$

Thus $T_{1212132}C_u \in \mathcal{M}$ as desired.

We prove (2) and (3). Let $w_1 = s_1s_2s_3s_2s_1s_2$ and $w_2 = s_2s_3s_2s_1s_2s_1$. These are both translations. Any element in $X_v$ can be written under the form $y_1w_1^n$ or $y_2w_2^n$ (where $y_1 \in Y(w_1)$ and $y_2 \in Y(w_2)$). Now we have

$$\begin{array}{rcll}
T_{s_2}C_v & \equiv & T_{s_2v} & \mod \mathcal{H}_{<0} \\
T_{s_1s_2}C_v & \equiv & T_{s_1s_2v} & \mod \mathcal{H}_{<0} \\
T_{s_2s_1s_2}C_v & \equiv & T_{s_2s_1s_2v} + T_{v_3} & \mod \mathcal{H}_{<0} \\
T_{s_3s_2s_1s_2}C_v & \equiv & T_{s_3s_2s_1s_2v} + T_{s_3v_3} & \mod \mathcal{H}_{<0} \\
T_{s_2s_3s_2s_1s_2}C_v & \equiv & T_{s_2s_3s_2s_1s_2v} + T_{s_2s_3v_3} & \mod \mathcal{H}_{<0} \\
T_{w_1}C_v & \equiv & T_{w_1v} + T_{s_1s_2s_3v_3} & \mod \mathcal{H}_{<0}
\end{array}$$

and if $T_z$ appears with a non zero coefficient in the expression of $T_{w_1}C_v$ in the standard basis then $\ell(w_1z) = \ell(w_1) + \ell(z)$. Therefore for all $y_1 \in Y(w_1)$ and $n \geq 1$ we have

$$T_{y_1w_1^n}C_v \equiv T_{yw_1^{n-1}s_2s_1s_2v} + T_{y_1w_1^{n-1}s_1s_2s_3v_3} \mod \mathcal{H}_{<0}.$$

Next we have

$$\begin{array}{rcll}
T_{s_2}C_v & \equiv & T_{s_2v} & \mod \mathcal{H}_{<0}\\
T_{s_3s_2}C_v & \equiv & T_{s_3s_2v} & \mod \mathcal{H}_{<0}\\
T_{s_1s_3s_2}C_v & \equiv & T_{s_1s_3s_2v} & \mod \mathcal{H}_{<0}\\
T_{s_2s_1s_3s_2}C_v & \equiv & T_{s_2s_1s_3s_2v} & \mod \mathcal{H}_{<0}\\
T_{s_1s_2s_1s_3s_2}C_v & \equiv & T_{s_1s_2s_1s_3s_2v} & \mod \mathcal{H}_{<0}\\
T_{s_2s_1s_2s_1s_3s_2}C_v & \equiv & T_{s_2s_1s_2s_1s_3s_2v} & \mod \mathcal{H}_{<0}\\
T_{s_3s_2s_1s_2s_1s_3s_2}C_v & \equiv & T_{s_3s_2s_1s_2s_1s_3s_2v} & \mod \mathcal{H}_{<0}\\
T_{w_2s_3s_2}C_v & \equiv & T_{w_2v} + T_{s_3s_2s_1s_2s_3v_3} & \mod \mathcal{H}_{<0}
\end{array}$$

and if $T_z$ appears with a non zero coefficient in the expression of $T_{w_2s_3s_2}C_v$ in the standard basis then $\ell(w_2z) = \ell(w_2) + \ell(z)$. Therefore for all $y_2 \in Y(w_2)$ and $n \geq 1$ we have

$$T_{y_2w_2^ns_3s_2}C_v \equiv T_{y_2w_2^{n-1}s_2s_1s_2v} + T_{y_2w_2^{n-1}s_3s_2s_1s_2s_3v_3} \mod \mathcal{H}_{<0}.$$

Let $y \in Y_v$ and let $y_0 = ys_2s_1s_2$. Using the previous equalities, we obtain that

$$T_yC_v \equiv T_{yv} + T_{y_0v_3} \mod \mathcal{H}_{<0}.$$

and for $x \in X_v - Y_v$ we have

$$T_yC_v \equiv T_{yv} \mod \mathcal{H}_{<0}.$$

Finally using the fact that $p^*_{xu,yv} \in \mathcal{A}_{<0}$ we get

$$\begin{array}{rcll}
\tilde{C_{yv}} & = & T_yC_v + \sum\limits_{\substack{u\in U, x\in X_u\\ xu\sqsubset yv}} p^*_{xu,yv}T_xC_u & \\
& = & T_yC_v + \sum\limits_{x<y, x\in X_v} p^*_{xv,yv}T_xC_v + \sum\limits_{\substack{u\in U, x\in X_u\\ u\neq v}} p^*_{xu,yv}T_xC_u & \\
& = & T_yC_v & \mod \mathcal{H}_{<0}\\
& = & T_{yv} + T_{y_0v_3} & \mod \mathcal{H}_{<0}
\end{array}$$

as required.

## 3. Proof of [3, Theorem 5.1] for $\tilde{G}_2$

We keep the notation of [3].

Let $(W, S)$ be the affine Weyl group of type $\tilde{G}_2$ with diagram and weight function given by



where $a, b$ are positive integers. The partition of $W$ into cells only depends on the ratio $r = a/b$. The aim of this section is to give some details in the computation involved in [3, Theorem 5.1]. Recall that in [3] we showed that the partition of $W$ given in [3, Conjecture 3.3] is constant whether

- $r > 2$
- $r = 2$
- $2 > r > 3/2$
- $r = 3/2$
- $3/2 > r > 1$
- $r = 1$
- $r < 1$

Fix $r > 0$. Let $\tilde{c}_0, \dots, \tilde{c}_n$ the sets obtained using the algorithm presented in [3, §3]. Then we denote by

- $\tilde{c}_i^j$ the left-connected components lying in $\tilde{c}_i$;
- $u_i^j$ the element of minimal length lying in $\tilde{c}_i^j$;
- $U$ the set which consists of all the $u_i^j$.

Let $u_i^j \in U$. We set

$$X_{u_i^j} := \{w \in W | wu_i^j \in \tilde{c}_i^j\}.$$

For each value of $r$ we give

1. the Hasse diagram of the pre-order $\preceq$ on $U$, which is also the left order on the left cells;
2. A table where for $u \in U$ and $x \in X_u$ we put the expression
$$T_{yv} + \sum_{xu \sqsubset yv} a_{xu,yv} T_x C_u$$
such that
$$T_y C_v \equiv T_{yv} + \sum_{xu \sqsubset yv} a_{xu,yv} T_x C_u \mod \mathcal{H}_{<0}.$$

*Remark* 3.1. When $T_x C_u \equiv T_{xu} \mod \mathcal{H}_{<0}$ we do not put it in the table.

For $w \in W$ we set

$$Y(w) = \{y \in W | \ell(wy^{-1}) = \ell(w) - \ell(y)\}.$$

- **Case $\mathbf{r > 2}$.**



FIGURE 2. Partition of $\tilde{G}_2$ into cells.

In the following Hasse diagram, the black edges hold for all $r > 2$, the green edges only hold for $3 < r \leq 4$ and the red edges only hold for $2 < r \leq 3$.



Figure 3. Order $\preceq$ on $U$

Let $w = s_1s_2s_1s_2s_3s_1s_2s_1s_2s_3$, $w_1 = s_1s_2s_3s_2s_1s_2$ and $w_2 = s_2s_3s_2s_1s_2s_1$. In the following table we have

$$y_1 \in \{yw_1^n | y \in Y(w_1), n \in \mathbb{N}\} \text{ and } y_2 \in \{yw_2^n | y \in Y(w_2)\}$$
$$y \in \{yw^n | y \in Y(w), n \in \mathbb{N}\}$$

TABLE 1. Condition **I5**

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_1^4$ | $s_1s_2s_1s_2s_3$ | $T_{xu} + v^{-a+3b}C_{u_0^6}$ |
| $u_1^4$ | $s_3s_1s_2s_1s_2s_3$ | $T_{xu} + v^{-a+3b}T_{s_3}C_{u_0^6}$ |
| $u_1^4$ | $s_2s_3s_1s_2s_1s_2s_3$ | $T_{xu} + v^{-a+3b}T_{s_2s_3}C_{u_0^6}$ |
| $u_1^4$ | $s_1s_2s_3s_1s_2s_1s_2s_3$ | $T_{xu} + v^{-a+3b}T_{s_1s_2s_3}C_{u_0^6}$ |
| $u_1^4$ | $s_2s_1s_2s_3s_1s_2s_1s_2s_3$ | $T_{xu} + v^{-a+3b}T_{s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_1^4$ | $yw$ | $T_{xu} + v^{-a+3b}T_yC_{u_0^6}$ |
| $u_1^6$ | $s_1s_2s_1s_2s_3$ | $T_{xu} + v^{-a+3b}C_{u_0^1}$ |
| $u_1^6$ | $s_3s_1s_2s_1s_2s_3$ | $T_{xu} + v^{-a+3b}T_{s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_2s_3s_1s_2s_1s_2s_3$ | $T_{xu} + v^{-a+3b}T_{s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_1s_2s_3s_1s_2s_1s_2s_3$ | $T_{xu} + v^{-a+3b}T_{s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_2s_1s_2s_3s_1s_2s_1s_2s_3$ | $T_{xu} + v^{-a+3b}T_{s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $yw$ | $T_{xu} + v^{-a+3b}T_yC_{u_0^1}$ |
| $u_2^3$ | $s_2s_1s_2$ | $T_{xu} + C_{u_0^6}$ |
| $u_2^3$ | $s_3s_2s_1s_2$ | $T_{xu} + T_{s_3}C_{u_0^6}$ |
| $u_2^3$ | $s_2s_3s_2s_1s_2$ | $T_{xu} + T_{s_2s_3}C_{u_0^6}$ |
| $u_2^3$ | $y_1w_1$ | $T_{xu} + T_{y_1s_1s_2s_3}C_{u_0^6}$ |
| $u_2^3$ | $y_2w_2s_3s_2$ | $T_{xu} + T_{y_2s_3s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_2^6$ | $s_2s_1s_2$ | $T_{xu} + C_{u_0^1}$ |
| $u_2^6$ | $s_3s_2s_1s_2$ | $T_{xu} + T_{s_3}C_{u_0^1}$ |
| $u_2^6$ | $s_2s_3s_2s_1s_2$ | $T_{xu} + T_{s_2s_3}C_{u_0^1}$ |
| $u_2^6$ | $y_1w_1$ | $T_{xu} + T_{y_1s_1s_2s_3}C_{u_0^1}$ |
| $u_2^6$ | $y_2w_2s_3s_2$ | $T_{xu} + T_{y_2s_3s_2s_1s_2s_3}C_{u_0^1}$ |

- **Case r = 2**



FIGURE 4. Partition of $\tilde{G}_2$ into cells



Figure 5. Order $\preceq$ on $U$

Let $w = s_1s_2s_1s_2s_3s_1s_2s_1s_2s_3$, $w_1 = s_1s_2s_3s_2s_1s_2$ and $w_2 = s_2s_3s_2s_1s_2s_1$. In the following table we have

$$y \in \{zw^n | z \in Y(w), n \in \mathbb{N}\}$$

$$y_1 \in \{yw_1^n | y \in Y(w_1), n \in \mathbb{N}\} \text{ and } y_2 \in \{yw_2^n | y \in Y(w_2)\}$$

Table 2. Condition **I5** for $r = 2$

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_1^2$ | $ys_1s_2s_1s_2s_3$ | $T_{xu} + T_yC_{u_0^1}$ |
| $u_1^4$ | $s_1s_2s_1s_2s_3$ | $T_{xu} + vC_{u_0^6} - C_{u_0^9}$ |
| $u_1^4$ | $s_3s_1s_2s_1s_2s_3$ | $T_{xu} + vT_{s_3}C_{u_0^6} - T_{s_3}C_{u_0^9}$ |
| $u_1^4$ | $s_2s_3s_1s_2s_1s_2s_3$ | $T_{xu} + vT_{s_2s_3}C_{u_0^6} - T_{s_2s_3}C_{u_0^9}$ |
| $u_1^4$ | $s_1s_2s_3s_1s_2s_1s_2s_3$ | $T_{xu} + vT_{s_1s_2s_3}C_{u_0^6} - T_{s_1s_2s_3}C_{u_0^9}$ |
| $u_1^4$ | $s_2s_1s_2s_3s_1s_2s_1s_2s_3$ | $T_{xu} + vT_{s_2s_1s_2s_3}C_{u_0^6} - T_{s_2s_1s_2s_3}C_{u_0^9}$ |
| $u_1^4$ | $yw$ | $T_{xu} + vT_{ys_1s_2s_1s_2s_3}C_{u_0^6} - T_{ys_2s_1s_2s_1s_2s_3}C_{u_0^6} - T_{ys_1s_2s_1s_2s_3}C_{u_0^9}$ |
| $u_1^6$ | $s_2s_1s_2s_1$ | $T_{xu} + vC_{u_0^1}$ |
| $u_1^6$ | $s_3s_1s_2s_1s_2s_3$ | $T_{xu} + vT_{s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_2s_3s_1s_2s_1s_2s_3$ | $T_{xu} + vT_{s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_1s_2s_3s_1s_2s_1s_2s_3$ | $T_{xu} + vT_{s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_2s_1s_2s_3s_1s_2s_1s_2s_3$ | $T_{xu} + vT_{s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $yw$ | $T_{xu} + vT_{ys_1s_2s_1s_2s_3}C_{u_0^1} - T_{ys_2s_1s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_2^3$ | $s_2s_1s_2$ | $T_{xu} + C_{u_0^6}$ |
| $u_2^3$ | $s_3s_2s_1s_2$ | $T_{xu} + T_{s_3}C_{u_0^6}$ |
| $u_2^3$ | $s_2s_3s_2s_1s_2$ | $T_{xu} + T_{s_2s_3}C_{u_0^6}$ |
| $u_2^3$ | $y_1w_1$ | $T_{xu} + T_{y_1s_1s_2s_3}C_{u_0^6}$ |
| $u_2^3$ | $y_2w_2s_3s_2$ | $T_{xu} + T_{y_2s_3s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_2^6$ | $s_2s_1s_2$ | $T_{xu} + C_{u_0^1}$ |
| $u_2^6$ | $s_3s_2s_1s_2$ | $T_{xu} + T_{s_3}C_{u_0^1}$ |
| $u_2^6$ | $s_2s_3s_2s_1s_2$ | $T_{xu} + T_{s_2s_3}C_{u_0^1}$ |
| $u_2^6$ | $y_1w_1$ | $T_{xu} + T_{y_1s_1s_2s_3}C_{u_0^1}$ |
| $u_2^6$ | $y_2w_2s_3s_2$ | $T_{xu} + T_{y_2s_3s_2s_1s_2s_3}C_{u_0^1}$ |

- **Case $\mathbf{3/2 < r < 2}$**



FIGURE 6. Partition of $\tilde{G}_2$ into cells



Figure 7. Order $\preceq$ on $U$

Let $w = s_1s_2s_1s_2s_3s_1s_2s_1s_2s_3$ and $w_1 = s_2s_3s_2s_1s_2s_1$. In the following table we have

$$y \in \{zw^n | z \in Y(w), n \in \mathbb{N}\} \text{ and } y_1 \in \{yw_1^n | y \in Y(w_1), n \in \mathbb{N}\}$$

## Table 3. Condition **I5** $3/2 < r < 2$

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_1^2$ | $ys_1s_2s_1s_2s_3$ | $T_{xu} + v^{-a+2b}T_yC_{u_0^1}$ |
| $u_1^4$ | $s_1s_2s_1s_2s_3$ | $T_{xu} + v^{-a+3b}C_{u_0^6} - v^{-a+2b}C_{u_0^9}$ |
| $u_1^4$ | $s_3s_1s_2s_1s_2s_3$ | $T_{xu} + v^{-a+3b}T_{s_3}C_{u_0^6} - v^{-a+2b}T_{s_3}C_{u_0^9}$ |
| $u_1^4$ | $s_2s_3s_1s_2s_1s_2s_3$ | $T_{xu} + v^{-a+3b}T_{s_2s_3}C_{u_0^6} - v^{-a+2b}T_{s_2s_3}C_{u_0^9}$ |
| $u_1^4$ | $s_1s_2s_3s_1s_2s_1s_2s_3$ | $T_{xu} + v^{-a+3b}T_{s_1s_2s_3}C_{u_0^6} - v^{-a+2b}T_{s_1s_2s_3}C_{u_0^9}$ |
| $u_1^4$ | $s_2s_1s_2s_3s_1s_2s_1s_2s_3$ | $T_{xu} + v^{-a+3b}T_{s_2s_1s_2s_3}C_{u_0^6} - v^{-a+2b}T_{s_2s_1s_2s_3}C_{u_0^9}$ |
| $u_1^4$ | $yw$ | $T_{xu} + v^{-a+3b}T_{ys_1s_2s_1s_2s_3}C_{u_0^6} - v^{-a+2b}T_{ys_2s_1s_2s_1s_2s_3}C_{u_0^6} - v^{-a+2b}T_{ys_1s_2s_1s_2s_3}C_{u_0^9}$ |
| $u_1^6$ | $s_2s_1s_2s_1$ | $T_{xu} + v^{-a+3b}C_{u_0^1}$ |
| $u_1^6$ | $s_3s_1s_2s_1s_2s_3$ | $T_{xu} + v^{-a+3b}T_{s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_2s_3s_1s_2s_1s_2s_3$ | $T_{xu} + v^{-a+3b}T_{s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_1s_2s_3s_1s_2s_1s_2s_3$ | $T_{xu} + v^{-a+3b}T_{s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_2s_1s_2s_3s_1s_2s_1s_2s_3$ | $T_{xu} + v^{-a+3b}T_{s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $yw$ | $T_{xu} + v^{-a+3b}T_{ys_1s_2s_1s_2s_3}C_{u_0^1} - v^{a-2b}T_{ys_2s_1s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_2^1$ | $s_2s_1s_2s_1$ | $T_{xu} + v^{a-b}C_{u_0^1}$ |
| $u_2^1$ | $s_3s_2s_1s_2s_1$ | $T_{xu} + v^{a-b}T_{s_2}C_{u_0^1}$ |
| $u_2^1$ | $y_1w_1$ | $T_{xu} + v^{a-b}T_{y_1s_2s_3}C_{u_0^1}$ |
| $u_2^1$ | $y_1w_1s_3s_2s_1$ | $T_{xu} + v^{a-b}T_{y_1s_3s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_2^3$ | $s_1s_2s_1$ | $T_{xu} - v^{a-b}C_{u_0^6}$ |
| $u_2^3$ | $s_2s_1s_2s_1$ | $T_{xu} - v^aC_{u_0^6}$ |
| $u_2^3$ | $s_3s_2s_1s_2s_1$ | $T_{xu} - v^aC_{s_3u_0^6}$ |
| $u_2^3$ | $y_1w_1$ | $T_{xu} - v^aT_{y_1s_3}C_{u_0^6} + T_{y_1s_3s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_2^3$ | $s_3s_2s_1$ | $T_{xu} - v^{a-b}T_{s_2s_3}C_{u_0^6}$ |
| $u_2^3$ | $s_1s_3s_2s_1$ | $T_{xu} - v^{a-b}T_{s_3}C_{u_0^6} + T_{s_1s_2s_3}C_{u_0^6}$ |
| $u_2^3$ | $s_2s_1s_3s_2s_1$ | $T_{xu} + v^{a+b}T_{s_2s_3}C_{u_0^6} + T_{s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_2^3$ | $s_1s_2s_1s_3s_2s_1$ | $T_{xu} + v^{a+b}T_{s_1s_2s_3}C_{u_0^6} + T_{s_1s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_2^3$ | $s_2s_1s_2s_1s_3s_2s_1$ | $T_{xu} + v^{a+b}T_{s_2s_1s_2s_3}C_{u_0^6} + T_{s_2s_1s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_2^3$ | $s_3s_2s_1s_2s_1s_3s_2s_1$ | $T_{xu} + v^{a+b}T_{s_3s_2s_1s_2s_3}C_{u_0^6} + T_{s_3s_2s_1s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_2^3$ | $y_1w_1s_3s_2s_1$ | $T_{xu} + v^aT_{y_1s_3s_2s_1s_2s_3}C_{u_0^6} + T_{y_1s_3s_2s_1s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_2^6$ | $s_1s_2s_1$ | $T_{xu} + v^{a+b}C_{u_0^1}$ |
| $u_2^6$ | $s_2s_1s_2s_1$ | $T_{xu} + v^aC_{u_0^1}$ |
| $u_2^6$ | $s_3s_2s_1s_2s_1$ | $T_{xu} + v^aC_{s_3u_0^1}$ |
| $u_2^6$ | $y_1w_1$ | $T_{xu} + v^aT_{y_1s_3}C_{u_0^1} + T_{y_1s_3s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_2^6$ | $s_3s_2s_1$ | $T_{xu} + v^{a+b}T_{s_2s_3}C_{u_0^1}$ |
| $u_2^6$ | $s_1s_3s_2s_1$ | $T_{xu} + v^{a+b}T_{s_3}C_{u_0^1} + T_{s_1s_2s_3}C_{u_0^1}$ |
| $u_2^6$ | $s_2s_1s_3s_2s_1$ | $T_{xu} + v^{a+b}T_{s_2s_3}C_{u_0^1} + T_{s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_2^6$ | $s_1s_2s_1s_3s_2s_1$ | $T_{xu} + v^{a+b}T_{s_1s_2s_3}C_{u_0^1} + T_{s_1s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_2^6$ | $s_2s_1s_2s_1s_3s_2s_1$ | $T_{xu} + v^{a+b}T_{s_2s_1s_2s_3}C_{u_0^1} + T_{s_2s_1s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_2^6$ | $s_3s_2s_1s_2s_1s_3s_2s_1$ | $T_{xu} + v^{a+b}T_{s_3s_2s_1s_2s_3}C_{u_0^1} + T_{s_3s_2s_1s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_2^6$ | $y_1w_1s_3s_2s_1$ | $T_{xu} + v^aT_{y_1s_3s_2s_1s_2s_3}C_{u_0^1} + T_{y_1s_3s_2s_1s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_3^3$ | $s_2s_1s_2$ | $T_{xu} + C_{u_0^6}$ |
| $u_3^3$ | $s_3s_2s_1s_2$ | $T_{xu} + T_{s_3}C_{u_0^6}$ |
| $u_3^5$ | $s_2s_1s_2$ | $T_{xu} + C_{u_0^1}$ |
| $u_3^5$ | $s_3s_2s_1s_2$ | $T_{xu} + T_{s_3}C_{u_0^1}$ |

- **Case $\mathbf{r = 3/2}$**



FIGURE 8. Partition of $\tilde{G}_2$ into cells



Figure 9. Order $\preceq$ on $U$

Let $w = s_1s_3s_2s_1s_2s_1s_3s_2s_1s_2$, and $w_1 = s_2s_3s_2s_1s_2s_1$. In the following table we have

$$y \in \{zw^n | z \in Y(w), n \in \mathbb{N}\} \text{ and } y_1 \in \{yw_1^n | y \in Y(w_1), n \in \mathbb{N}\}$$

TABLE 4. Condition **I5** for $r = 3/2$

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_1^1$ | $s_2s_1s_2s_1$ | $T_{xu} + vC_{u_0^1}$ |
| $u_1^1$ | $s_3s_2s_1s_2s_1$ | $T_{xu} + vT_{s_3}C_{u_0^1}$ |
| $u_1^1$ | $y_1w_1$ | $T_{xu} + vT_{ys_2s_3}C_{u_0^1}$ |
| $u_1^1$ | $y_1w_1s_3s_2s_1$ | $T_{xu} + vT_{ys_3s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_1^3$ | $s_1s_2s_1$ | $T_{xu} + vC_{u_0^6}$ |
| $u_1^3$ | $s_2s_1s_2s_1$ | $T_{xu} + v^3C_{u_0^6}$ |
| $u_1^3$ | $s_3s_2s_1s_2s_1$ | $T_{xu} + v^3T_{s_3}C_{u_0^6}$ |
| $u_1^3$ | $y_1w_1$ | $T_{xu} + v^3T_{ys_3}C_{u_0^6} + T_{ys_3s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_3s_2s_1$ | $T_{xu} + T_{s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_1s_3s_2s_1$ | $T_{xu} + T_{s_1s_2s_3}C_{u_0^6} + vT_{s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_2s_1s_2s_3}C_{u_0^6} + vT_{s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_1s_2s_1s_2s_3}C_{u_0^6} + vT_{s_1s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_2s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_2s_1s_2s_1s_2s_3}C_{u_0^6} + vT_{s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_3s_2s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_3s_2s_1s_2s_1s_2s_3}C_{u_0^6} + vT_{s_3s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $y_1w_1s_3s_2s_1$ | $T_{xu} + T_{ys_2s_3s_2s_1s_2s_1s_2s_3}C_{u_0^6} + v^3T_{ys_3s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_1^6$ | $s_1s_2s_1$ | $T_{xu} + vC_{u_0^1}$ |
| $u_1^6$ | $s_2s_1s_2s_1$ | $T_{xu} + v^3C_{u_0^1}$ |
| $u_1^6$ | $s_3s_2s_1s_2s_1$ | $T_{xu} + v^3T_{s_3}C_{u_0^1}$ |
| $u_1^6$ | $y_1w_1$ | $T_{xu} + v^3T_{ys_3}C_{u_0^1} + T_{ys_3s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_3s_2s_1$ | $T_{xu} + T_{s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_1s_3s_2s_1$ | $T_{xu} + T_{s_1s_2s_3}C_{u_0^1} + vT_{s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_2s_1s_2s_3}C_{u_0^1} + vT_{s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_1s_2s_1s_2s_3}C_{u_0^1} + vT_{s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_2s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_2s_1s_2s_1s_2s_3}C_{u_0^1} + T_{s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_3s_2s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_3s_2s_1s_2s_1s_2s_3}C_{u_0^1} + T_{s_3s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $y_1w_1s_3s_2s_1$ | $T_{xu} + T_{y_1s_2s_3s_2s_1s_2s_1s_2s_3}C_{u_0^1} + v^3T_{ys_3s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_2^2$ | $s_1s_3s_2s_1s_2$ | $T_{xu} + vT_{s_3}C_{u_0^1}$ |
| $u_2^2$ | $ys_1s_3s_2s_1s_2$ | $T_{xu} + vT_{ys_3}C_{u_0^1}$ |
| $u_2^4$ | $s_2s_1s_2$ | $T_{xu} + C_{u_0^6}$ |
| $u_2^4$ | $s_3s_2s_1s_2$ | $T_{xu} + T_{s_3}C_{u_0^6}$ |
| $u_2^4$ | $s_1s_3s_2s_1s_2$ | $T_{xu} + v^3T_{s_3}C_{u_0^6}$ |
| $u_2^4$ | $ys_1s_3s_2s_1s_2$ | $T_{xu} + v^3T_{ys_3}C_{u_0^6}$ |
| $u_2^6$ | $s_2s_1s_2$ | $T_{xu} + C_{u_0^1}$ |
| $u_2^6$ | $s_3s_2s_1s_2$ | $T_{xu} + T_{s_3}C_{u_0^1}$ |
| $u_2^6$ | $s_1s_3s_2s_1s_2$ | $T_{xu} + v^3T_{s_3}C_{u_0^1}$ |
| $u_2^6$ | $ys_1s_3s_2s_1s_2$ | $T_{xu} + v^3T_{ys_3}C_{u_0^1}$ |

- **Case $1 < r < 3/2$**



FIGURE 10. Partition of $\tilde{G}_2$ into cells



Figure 11. Order $\preceq$ on $U$

Let $w = s_1s_3s_2s_1s_2s_1s_3s_2s_1s_2$, and $w_1 = s_2s_3s_2s_1s_2s_1$. In the following table we have

$$y \in \{zw^n | z \in Y(w), n \in \mathbb{N}\} \text{ and } y_1 \in \{yw_1^n | y \in Y(w_1), n \in \mathbb{N}\}$$

TABLE 5. Condition **I5** for $1 < r < 3/2$

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_1^1$ | $s_2s_1s_2s_1$ | $T_{xu} + v^{a+b}C_{u_0^1}$ |
| $u_1^1$ | $s_3s_2s_1s_2s_1$ | $T_{xu} + v^{a+b}T_{s_3}C_{u_0^1}$ |
| $u_1^1$ | $y_1w_1$ | $T_{xu} + v^{a+b}T_{y_1s_2s_3}C_{u_0^1}$ |
| $u_1^1$ | $y_1w_1s_3s_2s_1$ | $T_{xu} + v^{a+b}T_{y_1s_3s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_1^3$ | $s_1s_2s_1$ | $T_{xu} + v^{a+b}C_{u_0^6}$ |
| $u_1^3$ | $s_2s_1s_2s_1$ | $T_{xu} + v^{a}C_{u_0^6}$ |
| $u_1^3$ | $s_3s_2s_1s_2s_1$ | $T_{xu} + v^{a}T_{s_3}C_{u_0^6}$ |
| $u_1^3$ | $y_1w_1$ | $T_{xu} + v^{a}T_{y_1s_2s_3}C_{u_0^6} + T_{y_1s_3s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_3s_2s_1$ | $T_{xu} + T_{s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_1s_3s_2s_1$ | $T_{xu} + T_{s_1s_2s_3}C_{u_0^6} + v^{a+b}T_{s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_2s_1s_2s_3}C_{u_0^6} + v^{a+b}T_{s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_1s_2s_1s_2s_3}C_{u_0^6} + v^{a+b}T_{s_1s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_2s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_2s_1s_2s_1s_2s_3}C_{u_0^6} + v^{a+b}T_{s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_3s_2s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_3s_2s_1s_2s_1s_2s_3}C_{u_0^6} + v^{a+b}T_{s_3s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $y_1w_1s_3s_2s_1$ | $T_{xu} + T_{y_1s_2s_3s_2s_1s_2s_1s_2s_3}C_{u_0^6} + v^{a}T_{ys_3s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_1^6$ | $s_1s_2s_1$ | $T_{xu} + v^{a+b}C_{u_0^1}$ |
| $u_1^6$ | $s_2s_1s_2s_1$ | $T_{xu} + v^{a}C_{u_0^1}$ |
| $u_1^6$ | $s_3s_2s_1s_2s_1$ | $T_{xu} + v^{a}T_{s_3}C_{u_0^1}$ |
| $u_1^6$ | $y_1w_1$ | $T_{xu} + v^{a}T_{ys_2s_3}C_{u_0^1} + T_{y_1s_3s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_3s_2s_1$ | $T_{xu} + T_{s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_1s_3s_2s_1$ | $T_{xu} + T_{s_1s_2s_3}C_{u_0^1} + v^{a+b}T_{s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_2s_1s_2s_3}C_{u_0^1} + v^{a+b}T_{s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_1s_2s_1s_2s_3}C_{u_0^1} + v^{a+b}T_{s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_2s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_2s_1s_2s_1s_2s_3}C_{u_0^1} + v^{a+b}T_{s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_3s_2s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_3s_2s_1s_2s_1s_2s_3}C_{u_0^1} + v^{a+b}T_{s_3s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $y_1w_1s_3s_2s_1$ | $T_{xu} + T_{y_1s_2s_3s_2s_1s_2s_1s_2s_3}C_{u_0^1} + v^{a}T_{y_1s_3s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_2^1$ | $s_1s_2$ | $T_{xu} + v^{a-b}C_{u_0^1}$ |
| $u_2^1$ | $s_2s_1s_2$ | $T_{xu} + v^{a}C_{u_0^1}$ |
| $u_2^1$ | $s_3s_2s_1s_2$ | $T_{xu} + v^{a}T_{s_3}C_{u_0^1}$ |
| $u_2^1$ | $s_1s_3s_2s_1s_2$ | $T_{xu} + (v^{2a} - v^{2a-2b})T_{s_3}C_{u_0^1}$ |
| $u_2^1$ | $ys_1s_3s_2s_1s_2$ | $T_{xu} + (v^{2a} - v^{2a-2b})T_{ys_3}C_{u_0^1}$ |
| $u_2^2$ | $s_1s_3s_2s_1s_2$ | $T_{xu} + v^{a+b}T_{s_3}C_{u_0^1}$ |
| $u_2^2$ | $ys_1s_3s_2s_1s_2$ | $T_{xu} + v^{a+b}T_{ys_3}C_{u_0^1}$ |
| $u_2^4$ | $s_2s_1s_2$ | $T_{xu} + C_{u_0^6}$ |
| $u_2^4$ | $s_3s_2s_1s_2$ | $T_{xu} + T_{s_3}C_{u_0^6}$ |
| $u_2^4$ | $s_1s_3s_2s_1s_2$ | $T_{xu} + v^{a}T_{s_3}C_{u_0^6}$ |
| $u_2^4$ | $ys_1s_3s_2s_1s_2$ | $T_{xu} + v^{a}T_{ys_3}C_{u_0^6}$ |
| $u_2^6$ | $s_2s_1s_2$ | $T_{xu} + C_{u_0^1}$ |
| $u_2^6$ | $s_3s_2s_1s_2$ | $T_{xu} + T_{s_3}C_{u_0^1}$ |
| $u_2^6$ | $s_1s_3s_2s_1s_2$ | $T_{xu} + v^{a}T_{s_3}C_{u_0^1}$ |
| $u_2^6$ | $ys_1s_3s_2s_1s_2$ | $T_{xu} + v^{a}T_{ys_3}C_{u_0^1}$ |

- **Case $\mathbf{r = 1}$**



FIGURE 12. Partition of $\tilde{G}_2$ into cells



Figure 13. Order $\preceq$ on $U$

Let $w = s_1s_3s_2s_1s_2s_1s_3s_2s_1s_2$, and $w_1 = s_2s_3s_2s_1s_2s_1$. In the following table we have

$$y \in \{zw^n | z \in Y(w), n \in \mathbb{N}\} \text{ and } y_1 \in \{yw_1^n | y \in Y(w_1), n \in \mathbb{N}\}$$

TABLE 6. Condition **I5** for $r = 1$

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_1^1$ | $s_2s_1s_2s_1$ | $T_{xu} + C_{u_0^1}$ |
| $u_1^1$ | $s_3s_2s_1s_2s_1$ | $T_{xu} + T_{s_3}C_{u_0^1}$ |
| $u_1^1$ | $y_1w_1$ | $T_{xu} + T_{y_1s_2s_3}C_{u_0^1}$ |
| $u_1^1$ | $y_1w_1s_3s_2s_1$ | $T_{xu} + T_{y_1s_3s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_1^3$ | $s_1s_2s_1$ | $T_{xu} + C_{u_0^6}$ |
| $u_1^3$ | $s_2s_1s_2s_1$ | $T_{xu} + vC_{u_0^6}$ |
| $u_1^3$ | $s_3s_2s_1s_2s_1$ | $T_{xu} + vT_{s_3}C_{u_0^6}$ |
| $u_1^3$ | $y_1w_1$ | $T_{xu} + vT_{y_1s_2s_3}C_{u_0^6} + T_{y_1s_3s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_3s_2s_1$ | $T_{xu} + T_{s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_1s_3s_2s_1$ | $T_{xu} + T_{s_1s_2s_3}C_{u_0^6} + T_{s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_2s_1s_2s_3}C_{u_0^6} + T_{s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_1s_2s_1s_2s_3}C_{u_0^6} + T_{s_1s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_2s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_2s_1s_2s_1s_2s_3}C_{u_0^6} + T_{s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_3s_2s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_3s_2s_1s_2s_1s_2s_3}C_{u_0^6} + T_{s_3s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $y_1w_1s_3s_2s_1$ | $T_{xu} + T_{y_1s_2s_3s_2s_1s_2s_1s_2s_3}C_{u_0^6} + T_{ys_3s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_1^6$ | $s_1s_2s_1$ | $T_{xu} + C_{u_0^1}$ |
| $u_1^6$ | $s_2s_1s_2s_1$ | $T_{xu} + vC_{u_0^1}$ |
| $u_1^6$ | $s_3s_2s_1s_2s_1$ | $T_{xu} + vT_{s_3}C_{u_0^1}$ |
| $u_1^6$ | $y_1w_1$ | $T_{xu} + vT_{y_1s_2s_3}C_{u_0^1} + T_{y_1s_3s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_3s_2s_1$ | $T_{xu} + T_{s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_1s_3s_2s_1$ | $T_{xu} + T_{s_1s_2s_3}C_{u_0^1} + T_{s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_2s_1s_2s_3}C_{u_0^1} + T_{s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_1s_2s_1s_2s_3}C_{u_0^1} + T_{s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_2s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_2s_1s_2s_1s_2s_3}C_{u_0^1} + T_{s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_3s_2s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_3s_2s_1s_2s_1s_2s_3}C_{u_0^1} + T_{s_3s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $y_1w_1s_3s_2s_1$ | $T_{xu} + T_{y_1s_2s_3s_2s_1s_2s_1s_2s_3}C_{u_0^1} + vT_{y_1s_3s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_2^1$ | $s_1s_2$ | $T_{xu} + C_{u_0^1}$ |
| $u_2^1$ | $s_2s_1s_2$ | $T_{xu} + vC_{u_0^1}$ |
| $u_2^1$ | $s_3s_2s_1s_2$ | $T_{xu} + vT_{s_3}C_{u_0^1}$ |
| $u_2^1$ | $s_1s_3s_2s_1s_2$ | $T_{xu} + v^2T_{s_3}C_{u_0^1}$ |
| $u_2^1$ | $ys_1s_3s_2s_1s_2$ | $T_{xu} + v^2T_{ys_3}C_{u_0^1}$ |
| $u_2^2$ | $ys_1s_3s_2s_1s_2$ | $T_{xu} + T_{ys_3}C_{u_0^1}$ |
| $u_2^4$ | $s_2s_1s_2$ | $T_{xu} + C_{u_0^6}$ |
| $u_2^4$ | $s_3s_2s_1s_2$ | $T_{xu} + T_{s_3}C_{u_0^6}$ |
| $u_2^4$ | $s_1s_3s_2s_1s_2$ | $T_{xu} + vT_{s_3}C_{u_0^6}$ |
| $u_2^4$ | $ys_1s_3s_2s_1s_2$ | $T_{xu} + vT_{ys_3}C_{u_0^6}$ |
| $u_2^6$ | $s_2s_1s_2$ | $T_{xu} + C_{u_0^1}$ |
| $u_2^6$ | $s_3s_2s_1s_2$ | $T_{xu} + T_{s_3}C_{u_0^1}$ |
| $u_2^6$ | $s_1s_3s_2s_1s_2$ | $T_{xu} + vT_{s_3}C_{u_0^1}$ |
| $u_2^6$ | $ys_1s_3s_2s_1s_2$ | $T_{xu} + vT_{ys_3}C_{u_0^1}$ |

- **Case $0 < r < 1$**



FIGURE 14. Partition of $\tilde{G}_2$ into cells

The green edges only hold for $1/2 \leq r < 1$.



Figure 15. Order $\preceq$ on $U$

Let $w = s_1s_3s_2s_1s_2s_1s_3s_2s_1s_2$, and $w_1 = s_2s_3s_2s_1s_2s_1$. In the following table we have

$$y \in \{zw^n | z \in Y(w), n \in \mathbb{N}\} \text{ and } y_1 \in \{yw_1^n | y \in Y(w_1), n \in \mathbb{N}\}$$

TABLE 7. Condition **I5** for $0 < r < 1$

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_1^3$ | $s_2s_1s_2s_1$ | $T_{xu} + v^aC_{u_0^6}$ |
| $u_1^3$ | $s_3s_2s_1s_2s_1$ | $T_{xu} + v^aT_{s_3}C_{u_0^6}$ |
| $u_1^3$ | $y_1w_1$ | $T_{xu} + v^aT_{y_1s_2s_3}C_{u_0^6} + T_{y_1s_3s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_3s_2s_1$ | $T_{xu} + T_{s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_1s_3s_2s_1$ | $T_{xu} + T_{s_1s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_1s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_2s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_2s_1s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $s_3s_2s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_3s_2s_1s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_1^3$ | $y_1w_1s_3s_2s_1$ | $T_{xu} + T_{y_1s_2s_3s_2s_1s_2s_1s_2s_3}C_{u_0^6} + v^aT_{y_1s_3s_2s_1s_2s_3}C_{u_0^6}$ |
| $u_1^6$ | $s_2s_1s_2s_1$ | $T_{xu} + v^aC_{u_0^1}$ |
| $u_1^6$ | $s_3s_2s_1s_2s_1$ | $T_{xu} + v^aT_{s_3}C_{u_0^1}$ |
| $u_1^6$ | $y_1w_1$ | $T_{xu} + v^aT_{y_1s_2s_3}C_{u_0^1} + T_{y_1s_3s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_3s_2s_1$ | $T_{xu} + T_{s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_1s_3s_2s_1$ | $T_{xu} + T_{s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_1s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_2s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_2s_1s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $s_3s_2s_1s_2s_1s_3s_2s_1$ | $T_{xu} + T_{s_3s_2s_1s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_1^6$ | $y_1w_1s_3s_2s_1$ | $T_{xu} + T_{y_1s_2s_3s_2s_1s_2s_1s_2s_3}C_{u_0^1} + v^aT_{y_1s_3s_2s_1s_2s_3}C_{u_0^1}$ |
| $u_2^4$ | $s_2s_1s_2$ | $T_{xu} + C_{u_0^6}$ |
| $u_2^4$ | $s_3s_2s_1s_2$ | $T_{xu} + T_{s_3}C_{u_0^6}$ |
| $u_2^4$ | $s_1s_3s_2s_1s_2$ | $T_{xu} + v^aT_{s_3}C_{u_0^6}$ |
| $u_2^4$ | $ys_1s_3s_2s_1s_2$ | $T_{xu} + v^aT_{ys_3}C_{u_0^6}$ |
| $u_2^6$ | $s_2s_1s_2$ | $T_{xu} + C_{u_0^1}$ |
| $u_2^6$ | $s_3s_2s_1s_2$ | $T_{xu} + T_{s_3}C_{u_0^1}$ |
| $u_2^6$ | $s_1s_3s_2s_1s_2$ | $T_{xu} + v^aT_{s_3}C_{u_0^1}$ |
| $u_2^6$ | $ys_1s_3s_2s_1s_2$ | $T_{xu} + v^aT_{ys_3}C_{u_0^1}$ |

## 4. PARTITION OF THE AFFINE WEYL GROUP $\tilde{B}_2$ INTO CELLS

Let $(W, S)$ be the affine Weyl group of type $\tilde{B}_2$ with diagram and weight function given by



where $a, b, c$ are positive integers. The partition of $W$ into cells only depend on the ratios $r_1 = a/b$ and $r_2 = c/b$. Note that by symmetry of diagram we may assume that $a \geq c$, that is $r_1 \geq r_2$. We define the following open subsets of $\mathbb{R}^2$:



*Remark* 4.1. Let $V$ be an Euclidean space of dimension 3 with standard basis $v_1, v_2, v_3$ corresponding respectively to the conjugacy classes $\{s_1\}$, $\{s_2\}$ and $\{s_3\}$ in $S$. In [3] we claimed that the essential hyperplanes of $W$ were

$$\mathcal{H}_{(1,0,0)}, \mathcal{H}_{(0,1,0)}, \mathcal{H}_{(0,0,1)}, \mathcal{H}_{(\epsilon,\epsilon,0)}, \mathcal{H}_{(0,\epsilon,\epsilon)}, \mathcal{H}_{(\epsilon,0,\epsilon)}, \mathcal{H}_{(\epsilon,\epsilon,\epsilon)}, \mathcal{H}_{(\epsilon,2\epsilon,\epsilon)}$$

where $\epsilon = \pm 1$. The picture above is just the projection of this hyperplane arrangement on the hyperplane $y = 1$.

4.1. **Partitions of $W$.** We describe the partitions $\mathcal{P}_{\mathcal{LR},\mathfrak{C}}$ and $\mathcal{P}_{\mathfrak{C}}$ of $\mathfrak{C}$ and the corresponding partition of $W$ according to [3, Conjecture 3.3]. We keep the notation of [3, §5].

- Case $r_1 = r_2, r_2 > 1$

Table 8. Partition $\mathcal{P}_{LR,\mathfrak{C}}$ and values of the $\mathbf{a}$-function

| | |
|---|---|
| $b_8 = \{e\}$ | $0$ |
| $b_7 = \{s_2\}$ | $b$ |
| $b_6 = \{s_1, s_1s_2, s_2s_1, s_2s_1s_2\}$ | $a$ |
| $b_5 = \{s_3, s_3s_2, s_2s_3, s_2s_3s_2\}$ | $a$ |
| $b_4 = \{s_3s_2s_3\}$ | $2a - b$ |
| $b_3 = \{s_1s_2s_1\}$ | $2a - b$ |
| $b_2 = \{s_1s_3\}$ | $2a$ |
| $b_1 = \{w_{2,3}\}$ | $2a + 2b$ |
| $b_0 = \{w_{1,2}\}$ | $2a + 2b$ |

Table 9. Partition $\mathcal{P}_{\mathfrak{C}}$

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| $r_2 > 1$ | $b_0 \cup b_1$ | $b_2$ | $b_3$ | $b_4$ | $b_5 \cup b_6$ | $b_7$ | $b_8$ |



$r_1 = r_2, r_2 > 1$

- Case $r_1 > 1$ and $r_2 > 1$

TABLE 10. Partition $\mathcal{P}_{LR,\mathfrak{C}}$ and values of the **a**-function

| | |
|---|---|
| $b_8 = \{e\}$ | $0$ |
| $b_7 = \{s_2\}$ | $b$ |
| $b_6 = \{s_3, s_2s_3, s_3s_2, s_2s_3s_2\}$ | $c$ |
| $b_5 = \{s_3s_2s_3\}$ | $2c - b$ |
| $b_4 = \{s_2s_3s_2s_3\}$ | $2b + 2c$ |
| $b_3 = \{s_1, s_2s_1, s_1s_2, s_2s_1s_2\}$ | $a$ |
| $b_2 = \{s_1s_3\}$ | $a + c$ |
| $b_1 = \{s_1s_2s_1\}$ | $2a - b$ |
| $b_0 = \{w_{1,2}\}$ | $2a + 2b$ |

TABLE 11. Partition $\mathcal{P}_{\mathfrak{C}}$

| | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| $(r_1, r_2) \in C_3$ | $b_0$ | $b_1$ | $b_2$ | $b_3 \leftrightarrow b_4$ | | $b_5$ | $b_6$ | $b_7$ | $b_8$ |
| $r_1 - r_2 - 2 = 0$ | $b_0$ | $b_1$ | $b_2 \cup b_4$ | | $b_3$ | $b_5$ | $b_6$ | $b_7$ | $b_8$ |
| $(r_1, r_2) \in C_2$ | $b_0$ | $b_1 \leftrightarrow b_4$ | | $b_2$ | $b_3$ | $b_5$ | $b_6$ | $b_7$ | $b_8$ |
| $r_1 - r_2 - 1 = 0$ | $b_0$ | $b_4$ | $b_1 \cup b_2$ | | $b_3$ | $b_5$ | $b_6$ | $b_7$ | $b_8$ |
| $(r_1, r_2) \in C_1$ | $b_0$ | $b_4$ | $b_2$ | $b_1$ | $b_3$ | $b_5$ | $b_6$ | $b_7$ | $b_8$ |


$(r_1, r_2) \in C_3$


$r_1 - r_2 - 2 = 0$



$(r_1, r_2) \in C_2$



$r_1 - r_2 - 1 = 0$



$(r_1, r_2) \in C_1$

- Case $r_1 > 1$ and $r_2 = 1$

TABLE 12. Partition $\mathcal{P}_{LR,\mathfrak{C}}$ and values of the **a**-function

| | |
|---|---|
| $b_6 = \{e\}$ | $0$ |
| $b_5 = \{s_2, s_3, s_3s_2, s_2s_3, s_3s_2s_3, s_2s_3s_2\}$ | $b$ |
| $b_4 = \{s_2s_3s_2s_3\}$ | $4b$ |
| $b_3 = \{s_1, s_2s_1, s_1s_2, s_2s_1s_2\}$ | $a$ |
| $b_2 = \{s_1s_3\}$ | $a + b$ |
| $b_1 = \{s_1s_2s_1\}$ | $2a - b$ |
| $b_0 = \{w_{1,2}\}$ | $2a + 2b$ |

Table 13. Partition $\mathcal{P}_{\mathfrak{C}}$

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| $r_1 > 3$ | $b_0$ | $b_1$ | $b_2$ | $b_3 \leftrightarrow b_4$ | | $b_5$ | $b_6$ |
| $r_1 = 3$ | $b_0$ | $b_1$ | $b_2 \cup b_4$ | | $b_3$ | $b_5$ | $b_6$ |
| $3 > r_1 > 2$ | $b_0$ | $b_1 \leftrightarrow b_4$ | | $b_2$ | $b_3$ | $b_5$ | $b_6$ |
| $r_1 = 2$ | $b_0$ | $b_4$ | $b_1 \cup b_2$ | | $b_3$ | $b_5$ | $b_6$ |
| $2 > r_1 > 1$ | $b_0$ | $b_4$ | $b_2$ | $b_1$ | $b_3$ | $b_5$ | $b_6$ |



$r_1 > 3$



$r_1 = 3$



$3 > r_1 > 2$



$r_1 = 2$



$2 > r_1 > 1$

- Case $r_1 > 1, r_2 < 1$

Table 14. Partition $\mathcal{P}_{LR,\mathfrak{c}}$ and values of the $\mathbf{a}$-function

| | |
|---|---|
| $b_8 = \{e\}$ | $0$ |
| $b_7 = \{s_3\}$ | $c$ |
| $b_6 = \{s_2, s_3s_2, s_2s_3, s_3s_2s_3\}$ | $b$ |
| $b_5 = \{s_2s_3s_2\}$ | $2b - c$ |
| $b_4 = \{s_2s_3s_2s_3\}$ | $2b + 2c$ |
| $b_3 = \{s_1, s_2s_1, s_1s_2, s_2s_1s_2\}$ | $a$ |
| $b_2 = \{s_1s_3\}$ | $a + c$ |
| $b_1 = \{s_1s_2s_1\}$ | $2a - b$ |
| $b_0 = \{w_{1,2}\}$ | $2a + 2b$ |

Table 15. Partition $\mathcal{P}_{\mathfrak{c}}$

| | |
|---|---|
| $(r_1, r_2) \in A_1$ | $b_0 \quad b_1 \quad b_2 \quad b_3 \leftrightarrow b_4 \quad b_5 \quad b_6 \quad b_7 \quad b_8$ |
| $r_1 - r_2 - 2 = 0$ | $b_0 \quad b_1 \quad b_2 \cup b_4 \quad b_3 \quad b_5 \quad b_6 \quad b_7 \quad b_8$ |
| $(r_1, r_2) \in A_2$ | $b_0 \quad b_1 \quad b_4 \quad b_2 \quad b_3 \quad b_5 \quad b_6 \quad b_7 \quad b_8$ |
| $r_1 + r_2 - 2 = 0, r_1 - r_2 > 1$ | $b_0 \quad b_1 \quad b_4 \quad b_2 \quad b_3 \cup b_5 \quad b_6 \quad b_7 \quad b_8$ |
| $(r_1, r_2) \in A_3$ | $b_0 \quad b_1 \leftrightarrow b_4 \quad b_2 \quad b_5 \quad b_3 \quad b_6 \quad b_7 \quad b_8$ |
| $r_1 - r_2 - 1 = 0, r_1 + r_2 < 2$ | $b_0 \quad b_4 \quad b_1 \cup b_2 \quad b_5 \quad b_3 \quad b_6 \quad b_7 \quad b_8$ |
| $(r_1, r_2) \in A_4$ | $b_0 \quad b_4 \quad b_2 \quad b_1 \quad b_5 \quad b_3 \quad b_6 \quad b_7 \quad b_8$ |
| $r_1 + r_2 - 2 = 0, r_1 - r_2 < 1$ | $b_0 \quad b_4 \quad b_2 \quad b_1 \quad b_5 \cup b_3 \quad b_6 \quad b_7 \quad b_8$ |
| $(r_1, r_2) \in A_5$ | $b_0 \quad b_4 \quad b_2 \quad b_1 \quad b_3 \quad b_5 \quad b_6 \quad b_7 \quad b_8$ |
| $r_1 - r_2 - 1 = 0, r_1 + r_2 > 2$ | $b_0 \quad b_4 \quad b_1 \cup b_2 \quad b_3 \quad b_5 \quad b_6 \quad b_7 \quad b_8$ |
| $(r_1, r_2) = (3/2, 1/2)$ | $b_0 \quad b_4 \quad b_1 \cup b_2 \quad b_3 \cup b_5 \quad b_6 \quad b_7 \quad b_8$ |


$(r_1, r_2) \in A_1$


$r_1 - r_2 - 2 = 0$



$(r_1, r_2) \in A_2$



$r_1 + r_2 - 2 = 0, r_1 - r_2 > 1$



$(r_1, r_2) \in A_3$



$r_1 + r_2 - 1 = 0, r_1 + r_2 > 2$



$(r_1, r_2) \in A_4$



$r_1 + r_2 - 2 = 0, r_1 - r_2 < 1$



$(r_1, r_2) \in A_5$



$r_1 - r_2 - 1 = 0, r_1 + r_2 > 2$



$(r_1, r_2) = (3/2, 1/2)$

- Case $r_1 = 1, r_2 < 1$

TABLE 16. Partition $\mathcal{P}_{LR,\mathfrak{C}}$ and values of the **a**-function

| | |
|---|---|
| $b_6 = \{e\}$ | $0$ |
| $b_5 = \{s_3\}$ | $c$ |
| $b_4 = \{s_2, s_1, s_1s_2, s_2s_1, s_1s_2s_1, s_3s_2, s_2s_3, s_3s_2s_3, s_2s_1s_2\}$ | $a$ |
| $b_3 = \{s_1s_3\}$ | $a + c$ |
| $b_2 = \{s_2s_3s_2\}$ | $2a - c$ |
| $b_1 = \{w_{2,3}\}$ | $2a + 2c$ |
| $b_0 = \{w_{1,2}\}$ | $4a$ |

TABLE 17. Partition $\mathcal{P}_{\mathfrak{C}}$

| | |
|---|---|
| $r_1 = 1, r_2 < 1$ | $b_0 \quad b_1 \quad b_2 \leftrightarrow b_3 \quad b_4 \quad b_5 \quad b_6$ |



$r_1 = 1, r_2 < 1$

- Case $r_1 < 1, r_2 < 1$

TABLE 18. Partition $\mathcal{P}_{LR,\mathfrak{C}}$ and values of the **a**-function

| | |
|---|---|
| $b_8 = \{e\}$ | $0$ |
| $b_7 = \{s_3\}$ | $c$ |
| $b_6 = \{s_1\}$ | $a$ |
| $b_5 = \{s_1s_3\}$ | $a + c$ |
| $b_4 = \{s_2, s_1s_2, s_2s_1, s_1s_2s_1, s_3s_2, s_2s_3, s_3s_2s_3\}$ | $b$ |
| $b_3 = \{s_2s_1s_2\}$ | $2b - a$ |
| $b_2 = \{s_2s_3s_2\}$ | $2b - c$ |
| $b_1 = \{w_{2,3}\}$ | $2b + 2c$ |
| $b_0 = \{w_{1,2}\}$ | $2a + 2b$ |

TABLE 19. Partition $\mathcal{P}_{\mathfrak{C}}$

| | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| $(r_1, r_2) \in B_2$ | $b_0$ | $b_1$ | $b_2$ | $b_3$ | $b_4$ | $b_5$ | $b_6$ | $b_7$ | $b_8$ |
| $r_1 + r_2 - 1 = 0$ | $b_0$ | $b_1$ | $b_2$ | $b_3$ | $b_4 \cup b_5$ | | $b_6$ | $b_7$ | $b_8$ |
| $(r_1, r_2) \in B_1$ | $b_0$ | $b_1$ | $b_2$ | $b_3 \leftrightarrow b_5$ | | $b_4$ | $b_6$ | $b_7$ | $b_8$ |


$(r_1, r_2) \in B_2$


$r_1 + r_2 - 1 = 0$


$(r_1, r_2) \in B_1$

- Case $r_1 = r_2, r_2 < 1$

Table 20. Partition $\mathcal{P}_{LR,\mathfrak{C}}$ and values of the $\mathbf{a}$-function

| | |
|---|---|
| $b_8 = \{e\}$ | $0$ |
| $b_7 = \{s_3\}$ | $a$ |
| $b_6 = \{s_1\}$ | $a$ |
| $b_5 = \{s_1s_3\}$ | $2a$ |
| $b_4 = \{s_2, s_1s_2, s_2s_1, s_1s_2s_1, s_3s_2, s_2s_3, s_3s_2s_3\}$ | $b$ |
| $b_3 = \{s_2s_3s_2\}$ | $2b - a$ |
| $b_2 = \{s_2s_1s_2\}$ | $2b - a$ |
| $b_1 = \{w_{2,3}\}$ | $2a + 2b$ |
| $b_0 = \{w_{1,2}\}$ | $2a + 2b$ |

Table 21. Partition $\mathcal{P}_{\mathfrak{C}}$

| | |
|---|---|
| $1 > r_2 > 1/2$ | $(b_0 \cup b_1) \quad (b_2 \cup b_3) \quad b_4 \quad b_5 \quad b_6 \quad b_7 \quad b_8$ |
| $r_2 = 1/2$ | $(b_0 \cup b_1) \quad (b_2 \cup b_3) \quad b_4 \cup b_5 \quad b_6 \quad b_7 \quad b_8$ |
| $r_2 < 1/2$ | $(b_0 \cup b_1) \quad (b_2 \cup b_3) \overset{\frac{3}{2}}{\leftrightarrow} b_5 \quad b_4 \quad b_6 \quad b_7 \quad b_8$ |



$1 > r_2 > 1/2$



$(r_1, r_2) = (1/2, 1/2)$



$1/2 > r_2 > 0$

- Case $r_1 = r_2 = 1$

TABLE 22. Partition $\mathcal{P}_{LR,\mathfrak{C}}$ and values of the **a**-function

| | |
|---|---|
| $b_4 = \{e\}$ | $0$ |
| $b_3 = \mathfrak{C} - b_0 \cup b_1 \cup \{e\}$ | $a$ |
| $b_2 = \{s_1 s_3\}$ | $2a$ |
| $b_1 = \{w_{2,3}\}$ | $4a$ |
| $b_0 = \{w_{1,2}\}$ | $4a$ |

TABLE 23. Partition $\mathcal{P}_{\mathfrak{C}}$

| | | | | |
|---|---|---|---|---|
| $r_2 = r_1 = 1$ | $b_0 \cup b_1$ | $b_2$ | $b_3$ | $b_4$ |



$r_1 = 1$ and $r_2 = 1$

4.2. **asymptotic cases.** We now consider the case where some parameters are equal to zero.

- Case $b = 0, a, c > 0$. We have

$$W = (\mathbb{Z}/2\mathbb{Z}) \ltimes \left(\tilde{A}_1 \times \tilde{A}_1\right)$$

where $\mathbb{Z}/2\mathbb{Z}$ is generated by $I = \{s_2\}$ and $\tilde{A}_1 \times \tilde{A}_1$ is generated by $s_1, s_2s_3s_2, s_3, s_2s_1s_2$. We know that the left (respectively two-sided) cells of $W$ are of the form $W_I.C$ (respectively $W_I.C.W_I$) where $C$ is a left (respectively two-sided) cell of $\tilde{A}_1 \times \tilde{A}_1$ with respect to the weight function $\tilde{L}$ defined by

$$\tilde{L}(s_1) = \tilde{L}(s_2s_1s_2) = a \text{ and } \tilde{L}(s_3) = \tilde{L}(s_2s_3s_2) = c.$$

We obtain the following partition of $W$.



$b = 0, a > c$



$b = 0, a = c$

- **Case** $c = 0$. We have

$$W = (\mathbb{Z}/2\mathbb{Z}) \ltimes \tilde{B}_2$$

where $(\mathbb{Z}/2\mathbb{Z})$ is generated by $s_3$ and $\tilde{B}_2$ is generated by $s_2, s_1, s_3s_2s_3$. We know that the left (respectively two-sided) cells of $W$ are of the form $W_I.C$ (respectively $W_I.C.W_I$) where $C$ is a left (respectively two-sided) cell of $\tilde{B}_2$ with respect to the weight function $\tilde{L}$ defined by

$$\tilde{L}(s_1) = a \text{ and } \tilde{L}(s_2) = \tilde{L}(s_3s_2s_3) = b.$$

We obtain the following partition of $W$.



$c = 0, b = 0, a > 0$



$c = 0, a > 2b$



$c = 0, a = 2b$



$c = 0, b < a < 2b$



$c = 0, a = b$



$c = 0, a < b$



$c = 0, a = 0, b > 0$

## 5. PROOF OF [3, Theorem 5.3] FOR $\tilde{B}_2$

As for $\tilde{G}_2$ (see Section 3), in each case we give the partial order $\preceq$ on $U$ (which is also the left order on the left cells) and a table for Condition **I5**. When we have $T_xC_u \equiv T_{xu}$ we do not put it in the table. So when there is no table it means that $T_xC_u \equiv T_{xu} \mod \mathcal{H}_{<0}$ for all $u \in U$ and $x \in X_u$.

- **Case** $(r_1, r_2) \in C_3$



FIGURE 16 Partition of $W$



FIGURE 17 Order $\preceq$ on $U$

- **Case** $r_1 - r_2 - 2 = 0$**,** $r_2 > 1$



FIGURE 18 Partition of $W$



FIGURE 19 Order $\preceq$ on $U$

- **Case** $(r_1, r_2) \in C_2$



FIGURE 20 Partition of $W$



FIGURE 21 Order $\preceq$ on $U$

- **Case** $r_1 - r_2 - 1 = 0$**,** $r_2 > 1$



FIGURE 22 Partition of $W$



FIGURE 23 Order $\preceq$ on $U$

- **Case** $(r_1, r_2) \in C_1$



FIGURE 24 Partition of $W$



FIGURE 25 Order $\preceq$ on $U$

Let $w = s_1s_3s_2s_1s_3s_2$ and $w_1 = s_1s_2s_3s_2$. In the following table

$$y \in \{zw^n | z \in Y(w), n \in \mathbb{N}\} \text{ and } y_1 \in \{zw^n | z \in Y(w_1), n \in \mathbb{N}\}.$$

TABLE 24. Condition **I5**

| $u \in U$ | $x \in X_u$ | $T_x C_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_2^1$ | $s_1 s_2$ | $T_{xu} + v^{a-c}\, C_{u_0^1}$ |
| $u_2^1$ | $y s_1 s_3 s_2$ | $T_{xu} + v^{a-c}\, T_{y s_3} C_{u_0^1}$ |
| $u_4^3$ | $s_2 s_3 s_2$ | $T_{xu} + v^{-a+2b}\, C_{u_1^3}$ |
| $u_4^3$ | $y_1 s_1 s_2 s_3 s_2$ | $T_{xu} + v^{-a+2b}\, T_{y_1 s_1} C_{u_1^3}$ |

- **Case** $r_1 = r_2, r_2 > 1$



Figure 26 Partition of $W$



Figure 27 Order $\preceq$ on $U$

Let $w = s_1s_3s_2s_1s_3s_2$. In the following table

$$y \in \{zw^n | y \in Y(w), n \in \mathbb{N}\}.$$

TABLE 25. Condition **I5**

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_2^1$ | $s_1s_2$ | $T_{xu} + C_{u_0^1}$ |
| $u_2^1$ | $ys_1s_3s_2$ | $T_{xu} + T_{ys_3}C_{u_0^1}$ |
| $u_2^3$ | $s_3s_2$ | $T_{xu} + C_{u_0^6}$ |
| $u_2^3$ | $ys_1s_3s_2$ | $T_{xu} + T_{ys_1}C_{u_0^6}$ |

- **Case** $r_1 > 3, r_2 = 1$



FIGURE 28 Partition of $W$



FIGURE 29 Order $\preceq$ on $U$

- **Case** $r_1 = 3, r_2 = 1$



FIGURE 30 Partition of $W$



FIGURE 31 Order $\preceq$ on $U$

- **Case** $2 < r_1 < 3, r_2 = 1$



FIGURE 32 Partition of $W$



FIGURE 33 Order $\preceq$ on $U$

- **Case** $r_1 = 2, r_2 = 1$



FIGURE 34 Partition of $W$



FIGURE 35 Order $\preceq$ on $U$

Let $w = s_1s_2s_3s_2$. In the following table

$$y \in \{zw^n | z \in Y(w), n \in \mathbb{N}\}.$$

Table 26. Condition **I5**

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_3^3$ | $s_2s_3s_2$ | $T_{xu} + C_{u_1^3}$ |
| $u_3^3$ | $ys_1s_2s_3s_2$ | $T_{xu} + T_{ys_1}C_{u_1^3}$ |

- **Case** $1 < r_1 < 2, r_2 = 1$



FIGURE 36 Partition of $W$



FIGURE 37 Order $\preceq$ on $U$

Let $w = s_1s_3s_2s_1s_3s_2$ and $w_1 = s_1s_2s_3s_2$. In the following table

$$y \in \{zw^n | z \in Y(w), n \in \mathbb{N}\} \text{ and } y_1 \in \{zw_1^n | z \in Y(w_1), n \in \mathbb{N}\}$$

Table 27. Condition **I5**

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_2^1$ | $s_1s_2$ | $T_{xu} + v^{a-c}\,C_{u_0^1}$ |
| $u_2^1$ | $ys_1s_3s_2$ | $T_{xu} + v^{a-c}\,T_{ys_3}C_{u_0^1}$ |
| $u_4^3$ | $s_2s_3s_2$ | $T_{xu} + v^{-a+2c}\,C_{u_1^3}$ |
| $u_4^3$ | $y_1w_1$ | $T_{xu} + v^{-a+2c}\,T_{ys_1}C_{u_1^3}$ |

- **Case** $(r_1, r_2) \in A_1$



FIGURE 38 Partition of $W$



FIGURE 39 Order $\preceq$ on $U$

- **Case** $r_1 - r_2 - 2 = 0$, $r_2 < 1$



FIGURE 40 Partition of $W$



FIGURE 41 Order $\preceq$ on $U$

- **Case** $(r_1, r_2) \in A_2$



FIGURE 42 Partition of $W$



FIGURE 43 Order $\preceq$ on $U$

Let $w_1 = s_1s_2s_3s_2$. In the following table

$$y_1 \in \{zw^n | z \in Y(w_1), n \in \mathbb{N}\}$$

TABLE 28. Condition **I5**

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_4^3$ | $s_2s_3s_2$ | $T_{xu} + v^{-a+2b}\, C_{u_2^3}$ |
| $u_4^3$ | $y_1w_1$ | $T_{xu} + v^{-a+2b}\, T_{y_1s_1}C_{u_2^3}$ |

- **Case** $r_1 + r_2 - 2 = 0, r_1 - r_2 > 1$



FIGURE 44 Partition of $W$



FIGURE 45 Order $\preceq$ on $U$

- **Case** $(r_1, r_2) \in A_3$



FIGURE 46 Partition of $W$



FIGURE 47 Order $\preceq$ on $U$

Let $w = s_2s_3s_2s_1$. In the following table

$$y \in \{zw^n | z \in Y(w), n \in \mathbb{N}\}.$$

Table 29. Condition **I5**

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_4^2$ | $s_2s_1$ | $T_{xu} - v^cC_{u_0^4}$ |
| $u_4^2$ | $s_3s_2s_1$ | $T_{xu} - v^cT_{s_3}C_{u_0^4}$ |
| $u_4^2$ | $ys_2s_3s_2s_1$ | $T_{xu} + T_{ys_3s_2s_3}C_{u_0^4} - v^cT_{ys_2s_3}C_{u_0^4}$ |
| $u_4^4$ | $s_2s_1$ | $T_{xu} - v^cC_{u_0^1}$ |
| $u_4^4$ | $s_3s_2s_1$ | $T_{xu} - v^cT_{s_3}C_{u_0^1}$ |
| $u_4^4$ | $ys_2s_3s_2s_1$ | $T_{xu} + T_{ys_3s_2s_3}C_{u_0^1} - v^cT_{ys_2s_3}C_{u_0^1}$ |

- **Case** $r_1 - r_2 - 1 = 0, r_1 + r_2 < 2$



FIGURE 48 Partition of $W$



FIGURE 49. Order $\preceq$ on $U$

Let $w = s_2s_3s_2s_1$. In the following table

$$y \in \{zw^n | z \in Y(w), n \in \mathbb{N}\}.$$

TABLE 30. Condition **I5**

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_3^2$ | $s_2s_1$ | $T_{xu} - v^cC_{u_0^4}$ |
| $u_3^2$ | $s_3s_2s_1$ | $T_{xu} - v^cT_{s_3}C_{u_0^4}$ |
| $u_3^2$ | $ys_2s_3s_2s_1$ | $T_{xu} + T_{ys_3s_2s_3}C_{u_0^4} - v^cT_{ys_2s_3}C_{u_0^4}$ |
| $u_3^4$ | $s_2s_1$ | $T_{xu} - v^cC_{u_0^1}$ |
| $u_3^4$ | $s_3s_2s_1$ | $T_{xu} - v^cT_{s_3}C_{u_0^1}$ |
| $u_3^4$ | $ys_2s_3s_2s_1$ | $T_{xu} + T_{ys_3s_2s_3}C_{u_0^1} - v^cT_{ys_2s_3}C_{u_0^1}$ |

- **Case** $(r_1, r_2) \in A_4$



FIGURE 50 Partition of $W$

In the following diagram, the green edges only hold for $a \geq 2c$.



FIGURE 51 Order $\preceq$ on $U$

Let $w = s_1s_3s_2s_1s_3s_2$ and $w_1 = s_2s_3s_2$. In the following table

$$y \in \{zw^n | z \in Y(w), n \in \mathbb{N}\} \text{ and } y_1 \in \{zw^n | z \in Y(w_1), n \in \mathbb{N}\}$$

Table 31. Condition **I5**

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_2^1$ | $s_1s_2$ | $T_{xu} + v^{a-c}C_{u_0^1}$ |
| $u_2^1$ | $ys_1s_3s_2$ | $T_{xu} + v^{a-c}T_{ys_3}C_{u_0^1}$ |
| $u_4^2$ | $s_2s_1$ | $T_{xu} - v^cC_{u_0^4}$ |
| $u_4^2$ | $s_3s_2s_1$ | $T_{xu} - v^cT_{s_3}C_{u_0^4}$ |
| $u_4^2$ | $y_1w_1$ | $T_{xu} + T_{y_1s_3s_2s_3}C_{u_0^4} - v^cT_{y_1s_2s_3}C_{u_0^4}$ |
| $u_4^4$ | $s_2s_1$ | $T_{xu} - v^cC_{u_0^1}$ |
| $u_4^4$ | $s_3s_2s_1$ | $T_{xu} - v^cT_{s_3}C_{u_0^1}$ |
| $u_4^4$ | $y_1w_1$ | $T_{xu} + T_{y_1s_3s_2s_3}C_{u_0^1} - v^cT_{y_1s_2s_3}C_{u_0^1}$ |

- **Case** $r_1 + r_2 - 2 = 0, r_1 - r_2 < 1$



FIGURE 52 Partition of $W$

In the following diagram, the green edges only hold for $a \geq 2c$.



FIGURE 53 Order $\preceq$ on $U$

Let $w = s_1s_3s_2s_1s_3s_2$. In the following table

$$y \in \{zw^n | z \in Y(w), n \in \mathbb{N}\}.$$

Table 32. Condition **I5**

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_2^1$ | $s_1s_2$ | $T_{xu} + v^{a-c}\, C_{u_0^1}$ |
| $u_2^1$ | $ys_1s_3s_2$ | $T_{xu} + v^{a-c}\, T_{ys_3}C_{u_0^1}$ |

- **Case** $(r_1, r_2) \in A_5$



FIGURE 54 Partition of $W$

In the following diagram, the green edges only hold for $a \geq 2c$.



FIGURE 55 Order $\preceq$ on $U$

Let $w = s_1s_3s_2s_1s_3s_2$ and $w_1 = s_1s_2s_3s_2$. In the following table

$$y \in \{zw^n | z \in Y(w), n \in \mathbb{N}\} \text{ and } y_1 \in \{zw_1^n | z \in Y(w_1), n \in \mathbb{N}\}$$

TABLE 33. Condition **I5**

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_2^1$ | $s_1s_2$ | $T_{xu} + v^{a-c}\, C_{u_0^1}$ |
| $u_2^1$ | $ys_1s_3s_2$ | $T_{xu} + v^{a-c}\, T_{ys_3}C_{u_0^1}$ |
| $u_4^3$ | $s_2s_3s_2$ | $T_{xu} + v^{-a+2b}\, C_{u_1^3}$ |
| $u_4^3$ | $y_1w_1$ | $T_{xu} + v^{-a+2b}\, T_{y_1s_1}C_{u_1^3}$ |

- **Case** $r_1 - r_2 - 1 = 0, r_1 + r_2 > 2$**,** $r_2 < 1$



Figure 56 Partition of $W$



Figure 57 Order $\preceq$ on $U$

Let $w_1 = s_1s_2s_3s_2$. In the following table

$$y_1 \in \{zw_1^n | z \in Y(w_1), n \in \mathbb{N}\}$$

TABLE 34. Condition **I5**

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_3^3$ | $s_2s_3s_2$ | $T_{xu} + v^{b-c}\, C_{u_1^3}$ |
| $u_3^3$ | $y_1s_1s_2s_3s_2$ | $T_{xu} + v^{b-c}\, T_{y_1s_1}C_{u_1^3}$ |

- **Case** $(r_1, r_2) = (3/2, 1/2)$



FIGURE 58 Partition of $W$



FIGURE 59 Order $\preceq$ on $U$

- **Case** $r_1 = 1, r_2 < 1$



FIGURE 60 Partition of $W$

In the following diagram, the green edges only hold for $a \geq 2c$.



FIGURE 61 Order $\preceq$ on $U$

Let $w = s_1s_3s_2s_1s_3s_2$ and $w_1 = s_2s_3s_2s_1$. In the following table

$$y \in \{zw^n | z \in Y(w), n \in \mathbb{N}\} \text{ and } y_1 \in \{zw^n | z \in Y(w_1), n \in \mathbb{N}\}$$

TABLE 35. Condition **I5**

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_2^2$ | $s_2s_1$ | $T_{xu} - v^cC_{u_0^4}$ |
| $u_2^2$ | $s_3s_2s_1$ | $T_{xu} - v^cT_{s_3}C_{u_0^4}$ |
| $u_2^2$ | $y_1w_1$ | $T_{xu} + T_{y_1s_3s_2s_3}C_{u_0^4} - v^cT_{y_1s_2s_3}C_{u_0^4}$ |
| $u_2^4$ | $s_2s_1$ | $T_{xu} - v^cC_{u_0^1}$ |
| $u_2^4$ | $s_3s_2s_1$ | $T_{xu} - v^cT_{s_3}C_{u_0^1}$ |
| $u_2^4$ | $ys_2s_3s_2s_1$ | $T_{xu} + T_{ys_3s_2s_3}C_{u_0^1} - v^cT_{ys_2s_3}C_{u_0^1}$ |
| $u_3^1$ | $s_1s_2$ | $T_{xu} + v^{a-c}C_{u_0^1}$ |
| $u_3^1$ | $ys_1s_3s_2$ | $T_{xu} + v^{a-c}T_{ys_3}C_{u_0^1}$ |

- Case $(r_1, r_2) \in B_1$



FIGURE 62 Partition of $W$

In the following diagram, the green edges only hold for $a \geq 2c$.



FIGURE 63 Order $\preceq$ on $U$

Let $w = s_1s_3s_2s_1s_3s_2$ and $w_1 = s_2s_3s_2s_1$. In the following table

$$y \in \{zw^n | z \in Y(w), n \in \mathbb{N}\} \text{ and } y_1 \in \{zw^n | z \in Y(w_1), n \in \mathbb{N}\}.$$

Table 36. Condition **I5**

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_2^2$ | $s_2s_1$ | $T_{xu} - v^cC_{u_0^4}$ |
| $u_2^2$ | $s_3s_2s_1$ | $T_{xu} - v^cT_{s_3}C_{u_0^4}$ |
| $u_2^2$ | $y_1w_1$ | $T_{xu} + T_{y_1s_3s_2s_3}C_{u_0^4} - v^cT_{y_1s_2s_3}C_{u_0^4}$ |
| $u_2^4$ | $s_2s_1$ | $T_{xu} - v^cC_{u_0^1}$ |
| $u_2^4$ | $s_3s_2s_1$ | $T_{xu} - v^cT_{s_3}C_{u_0^1}$ |
| $u_2^4$ | $y_1w_1$ | $T_{xu} + T_{y_1s_3s_2s_3}C_{u_0^1} - v^cT_{y_1s_2s_3}C_{u_0^1}$ |
| $u_3^1$ | $s_1s_2$ | $T_{xu} + v^{a-c}C_{u_0^1}$ |
| $u_3^1$ | $s_1s_3s_2$ | $T_{xu} + v^{a-c}T_{s_3}C_{u_0^1}$ |
| $u_3^1$ | $s_2s_1s_3s_2$ | $T_{xu} + v^{a-c}T_{s_2s_3}C_{u_0^1}$ |
| $u_3^1$ | $s_3s_2s_1s_3s_2$ | $T_{xu} + v^{a-c}T_{s_3s_2s_3}C_{u_0^1}$ |
| $u_3^1$ | $s_1s_2s_1s_3s_2$ | $T_{xu} + v^{a-c}T_{s_1s_2s_3}C_{u_0^1}$ |
| $u_3^1$ | $ys_1s_3s_2s_1s_3s_2$ | $T_{xu} + v^{a-c}T_{ys_1s_3s_2s_3}C_{u_0^1}$ |

- **Case** $r_1 + r_2 - 1 = 0$



FIGURE 64 Partition of $W$

In the following diagram, the green edges only hold for $a \geq 2c$.



FIGURE 65 Order $\preceq$ on $U$

Let $w_1 = s_2s_3s_2s_1$. In the following table

$$y_1 \in \{zw^n | z \in Y(w_1), n \in \mathbb{N}\}.$$

TABLE 37. Condition **I5**

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_2^2$ | $s_2s_1$ | $T_{xu} - v^cC_{u_0^4}$ |
| $u_2^2$ | $s_3s_2s_1$ | $T_{xu} - v^cT_{s_3}C_{u_0^4}$ |
| $u_2^2$ | $ys_2s_3s_2s_1$ | $T_{xu} + T_{ys_3s_2s_3}C_{u_0^4} - v^cT_{ys_2s_3}C_{u_0^4}$ |
| $u_2^4$ | $s_2s_1$ | $T_{xu} - v^cC_{u_0^1}$ |
| $u_2^4$ | $s_3s_2s_1$ | $T_{xu} - v^cT_{s_3}C_{u_0^1}$ |
| $u_2^4$ | $ys_2s_3s_2s_1$ | $T_{xu} + T_{ys_3s_2s_3}C_{u_0^1} - v^cT_{ys_2s_3}C_{u_0^1}$ |

- **Case** $(r_1, r_2) \in B_2$



FIGURE 66 Partition of $W$

In the following diagram, the green edges only hold for $a \geq 2c$.



FIGURE 67 Order $\preceq$ on $U$

Let $w_1 = s_2s_3s_2s_1$. In the following table

$$y_1 \in \{zw_1^n | z \in Y(w_1), n \in \mathbb{N}\}.$$

Table 38. Condition **I5**

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_2^2$ | $s_2s_1$ | $T_{xu} - v^cC_{u_0^4}$ |
| $u_2^2$ | $s_3s_2s_1$ | $T_{xu} - v^cT_{s_3}C_{u_0^4}$ |
| $u_2^2$ | $ys_2s_3s_2s_1$ | $T_{xu} + T_{ys_3s_2s_3}C_{u_0^4} - v^cT_{ys_2s_3}C_{u_0^4}$ |
| $u_2^4$ | $s_2s_1$ | $T_{xu} - v^cC_{u_0^1}$ |
| $u_2^4$ | $s_3s_2s_1$ | $T_{xu} - v^cT_{s_3}C_{u_0^1}$ |
| $u_2^4$ | $ys_2s_3s_2s_1$ | $T_{xu} + T_{ys_3s_2s_3}C_{u_0^1} - v^cT_{ys_2s_3}C_{u_0^1}$ |

- **Case** $r_1 = r_2, 1/2 < r_2 < 1$



FIGURE 68 Partition of $W$



FIGURE 69 Order $\preceq$ on $U$

Let $w = s_1s_3s_2s_1s_3s_2$. In the following table

$$y \in \{zw^n | z \in Y(w), n \in \mathbb{N}\}.$$

TABLE 39. Condition **I5**

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_2^1$ | $s_1s_2$ | $T_{xu} - +C_{u_0^1}$ |
| $u_2^1$ | $ys_1s_3s_2$ | $T_{xu} + T_{ys_3}C_{u_0^1}$ |
| $u_2^3$ | $s_3s_2$ | $T_{xu} + C_{u_0^6}$ |
| $u_2^3$ | $ys_1s_3s_2$ | $T_{xu} + T_{ys_1}C_{u_0^6}$ |

- **Case** $r_1 = r_2, r_2 = 1/2$



FIGURE 70 Partition of $W$



FIGURE 71 Order $\preceq$ on $U$

- **Case** $r_1 = r_2, 0 < r_2 < 1/2$



FIGURE 72 Partition of $W$



FIGURE 73 Order $\preceq$ on $U$

- **Case** $r_1 = r_2 = 1$



FIGURE 74 Partition of $W$



FIGURE 75 Order $\preceq$ on $U$

Let $w = s_1s_3s_2s_1s_3s_2$. In the following table

$$y \in \{zw^n | z \in Y(w), n \in \mathbb{N}\}.$$

TABLE 40. Condition **I5**

| $u \in U$ | $x \in X_u$ | $T_xC_u \mod \mathcal{H}_{<0}$ |
|---|---|---|
| $u_2^1$ | $s_1s_2$ | $T_{xu} + C_{u_0^1}$ |
| $u_2^1$ | $ys_1s_3s_2$ | $T_{xu} + T_{ys_3}C_{u_0^1}$ |
| $u_2^3$ | $s_3s_2$ | $T_{xu} + C_{u_0^6}$ |
| $u_2^3$ | $ys_1s_3s_2$ | $T_{xu} + T_{ys_1}C_{u_0^6}$ |

## REFERENCES

[1] J. Guilhot. On the determination of Kazhdan-Lusztig cells for affine Weyl group with unequal parameter. *J. of Algebra* **318**, *893–917*, 2007.

[2] J. Guilhot. Generalized induction of Kazhdan-Lusztig cells. *To appear in Ann. Inst. Fourier, available at http://arxiv.org/abs/math/0703764.*

[3] J. Guilhot. Kazhdan-lusztig cells in affine weyl groups of rank 2. *Preprint available at http://arxiv.org/abs/0901.1711.*

[4] Martin Schönert et al. *GAP – Groups, Algorithms, and Programming – version 3 release 4 patchlevel 4*. Lehrstuhl D für Mathematik, Rheinisch Westfälische Technische Hochschule, Aachen, Germany, 1997.

SCHOOL OF MATHEMATICS AND STATISTICS F07 UNIVERSITY OF SYDNEY NSW 2006 AUSTRALIA

*E-mail address*: `guilhot@maths.usyd.edu.au`